

The dark matter research

Steve Branchu

2 June 2017

Contents

1	Introduction	1
2	OHe atomic structure	1
3	Signatures of OHe	1
3.1	Positron annihilation in galaxy bubbles	1
3.2	OHe de-excitation	2
3.3	The LHC probes	2
4	Some potential problems	2
4.1	Inelastic process	2
4.2	Problems of OBe dark matter	5
5	Conclusion	5
6	References	6

1 Introduction

The dark matter, which corresponds to 25 % of total cosmologic density, makes it possible to understand the creation of the big structures in the Big Bang. This matter is named like this because it does not emit rays. Indeed its existence is deduced by the observation of the speed of rotations of the galaxies, which are not the same to values obtained by the luminosity law. The non baryonic nature of the dark matter comes from analysis of the nucleosynthesis of the Big Bang and the cosmic microwave background anisotropies.

The problem is that simulations of evolution of the Universe give different results according to the nature of the dark matter. To remove this ambiguity, astrophysicists try to determine the composition of this dark matter. Recent searches indicate that a combination between nucleus of He and 2 leptons, named OHe, possibly satisfy the characteristic of the dark matter.

The O-Helium dark matter is named composite dark matter model. I will explain in this report the research which permit to discover it, the advantages of this model and finally the problems of this model.

2 OHe atomic structure

An OHe atom is composed by a helium nucleus and an heavy double charged particle O^{--} (reference [1]), maybe a lepton. Here are the mass M_O and M_{He} , and the charge $Z_O = 2e$. The potential interaction follows the laws below :

$$V(r \geq R_{He}) = -\frac{Z_O Z_{He} \alpha}{r}$$
$$V(r \leq R_{He}) = -\frac{Z_O Z_{He} \alpha}{r} \left(3 - \frac{r^2}{R_{He}^2} \right)$$

where α is the fine structure constant and r the distance between the leptons and the center of the nucleus.

3 Signatures of OHe

3.1 Positron annihilation in galaxy bubbles

The satellite INTEGRAL observed a positron-annihilation line excess in the galactic bulge, and the number of collisions increases in the high gas density zone. It can be explained by the presence of OHe (references [1, 2]). The dark matter is more concentrated in the bulge than the rest of the galaxy, and this gas is collisionless. The collisions excite the OHe gas in creating a pair production, then the de-excitation in E0 transitions had

been observed by excess of positron-annihilation line. However, calculated rate of collisions depends of the density of OHe, but the theoretical estimations for dark matter and baryonic are uncertain. The last analysis indicates a little value of dark matter density, and deduce the mass of O^{--} near 1.0 TeV.

3.2 OHe de-excitation

The collisions excite the OHe gas at the first level with non-zero angular momentum, the E1 transition create gamma lines with energies :

$$E_{nm} = \left(\frac{1}{m^2} - \frac{1}{n^2} \right) \times 1.5879 \text{ MeV}$$

In all line emissions from the Galaxy (references [1]), there are some X-Ray and gamma ray, probably originates from OHe. The INTEGRAL and XMM-Newton data permit to compare the observations and the simulations to understand the unidentified lines.

3.3 The LHC probes

To understand how constitutes the OHe, so the nature of O^{--} , scientist at LHC done experiences with massive stable charged particles around 1 TeV (references [2]). The detector ATLAS and CMS detected particles around 700 GeV, so this is a good way to find the particle which correspond to composite dark matter.

4 Some potential problems

4.1 Inelastic process

During the young age of the Universe, inelastic diffusion between particles of OHe and between OHe and primordial He decreased the quantity of OHe (references [3]). The nuclear reactions are :



where Be is the beryllium. The OHe forms at a temperature T_0 which depends of its binding energy, which is 1.175 MeV, that corresponds to $T_0 = 50 \text{ keV}$. The cosmological time is calculated with the temperature, the inelastic process start at a time :

$$t(s) \simeq \frac{1}{T^2(\text{MeV})} \simeq \frac{1}{0.05^2} = 400s$$

after the Big Bang and continue until helium freezes out at $t_* \simeq 600s$.

During these 200 s, the **diminution** rate of the quantity of OHe is :

$$\frac{dn_{OHe}}{dt} = -3Hn_{OHe} - n_{OHe}^2\sigma_1v_1 - n_{OHe}n_{He}\sigma_2v_2 \quad (3)$$

with n_{OHe} and n_{He} are the densities of OHe and He, $H = 1/2t$ the expansion rate of the Universe during the radiation dominance era, σ_1 , σ_2 , v_1 and v_2 are respectively the cross section and the relative velocity of the reaction (1) and (2). The only factor which affect n_{He} is the expansion :

$$\frac{dn_{He}}{dt} = -3Hn_{He} \iff n_{He}(t) = n_{He}^0 \left(\frac{t_0}{t}\right)^{3/2} \quad (4)$$

where n_{He}^0 is the number density of He at $t = t_0$. We calculate the fraction of free OHe atoms due to their inelastic reactions, with the ratio f of the number density of OHe and He nuclei, $f = \frac{n_{OHe}}{n_{He}}$. The expressions (3) and (4) give the relation of evolution :

$$\frac{df}{dt} = -n_{He}f(\sigma_1v_1f + \sigma_2v_2) \quad (5)$$

The cross sections σ_1 and σ_2 depends of the size of the nucleus :

$$\sigma_1 \approx 4\pi(2r_{OHe})^2 \quad (6)$$

$$\sigma_2 \approx 4\pi(r_{OHe} + r_{He})^2 \quad (7)$$

where r are the radius of respectively the OHe nucleus and the He nucleus. The both are approximately equal to 2 fm, $\sigma_1 \approx \sigma_2 \approx 64\pi 10^{-26} cm^2$. The mean relative velocities v_1 and v_2 are obtained by the Maxwell-Boltzmann velocity distributions, because the species are in thermal equilibrium with the plasma :

$$v_1 = \sqrt{\frac{8T}{\pi\mu_1}} \quad (8)$$

$$v_2 = \sqrt{\frac{8T}{\pi\mu_2}} \quad (9)$$

with $\mu_1 = \frac{m_{OHe}}{2}$ the reduced mass of OHe-OHe and $\mu_2 \simeq m_{He}$ the reduced mass of OHe-He. During the radiation-dominated era, the relation between time and temperature is : $Tt^{1/2} = T_0t_0^{1/2}$, we insert it in the expression of velocities, and use the relation (5) to obtain :

$$\frac{df}{dt} = -\gamma \frac{1}{t^{7/4}} f(\alpha f + \beta) \quad (10)$$

with

$$\alpha = \frac{\sigma_1}{\sqrt{\mu_1}} \quad (11)$$

$$\beta = \frac{\sigma_2}{\sqrt{\mu_2}} \quad (12)$$

$$\gamma = n_{He}^0 t_0^{7/4} \sqrt{\frac{8T_0}{\pi}} \quad (13)$$

The initial condition is $f(t_0) = f_0$, the solution of (10) is :

$$f(t) = \frac{\beta f_0}{\exp(\frac{4}{3}\beta\gamma(t_0^{-3/4} - t^{-3/4}))(\alpha f_0 + \beta) - \alpha} \quad (14)$$

We can deduce the number density of He at the beginning of the **baryon** nucleosynthesis from its actual value. It corresponds today about 10% of all baryons, with a present critical density measured is to :

$$\rho_c^1 = 5.67 \times 10^{-6} m_p / cm^3 \iff n_{He}^1 \simeq 2.8 \times 10^{-8} cm^{-3} \quad (15)$$

Then, we have to add the effect of the expansion, which is proportionnal at the temperature of the CMB $n_{He}^0 = n_{He}^1 \left(\frac{T_0}{T_1}\right)^3 \simeq 2.8 \times 10^{-17} cm^{-3}$. At the **beginning** of the Universe the quantity of O^{--} **particules** was the same that OHe **particules**. Today the quantity of **particules** hasn't been change, but the size of the Universe increase so the density of particules has been diluted. The fraction of OHe hasn't been change, almost 25% of the critical density, so we can calculate the parameter $f_0 \simeq 0.05$. Now we can insert this value in the equation (14) and calculate the fraction at the freeze-out time :

$$f(t_*) \simeq 5 \times 10^{-6133} \ll f_0 \quad (16)$$

this results means the OHe reactions **don't during**  to be more clearly, the reaction (2) is majoritary, so the dark matter is now in form of OBe. **The suppression instant arrive when the exponential term has the value e^{14127} .** This value represents the number N_2 of

reactions that happened between t_0 and t_* :

$$\begin{aligned}
 N_2 &= \int_{t_0}^{t_*} n_{He}(t) \sigma_2 v_2(t) dt \\
 &= n_{He}^0 t_0^{3/2} \sigma_2 \sqrt{\frac{8T_0 t_0^{1/2}}{\pi \mu_2}} \int_{t_0}^{t_*} \frac{1}{t^{7/4}} dt \\
 &= n_{He}^0 t_0^{7/4} \sqrt{\frac{8T_0}{\pi}} \frac{\sigma_2}{\sqrt{\mu_2}} \left(-\frac{4}{3} \right) \left(\frac{1}{t_*^{3/4}} - \frac{1}{t_0^{3/4}} \right) \\
 &= \frac{4}{3} \beta \gamma \left(\frac{1}{t_0^{3/4}} - \frac{1}{t_*^{3/4}} \right)
 \end{aligned}$$

This relation means the realization of the OHe dark matter model implies a very strong suppression of the reaction (2), which corresponding to $N_2 \ll 1$. This particular case needs the development of a strong dipole Coulomb barrier in OHe-He interaction his one of the most important problem in this scenario.

4.2 Problems of OBe dark matter

Due to the Coulomb barrier, the particule of OBe **can't capture leptons**,  it recombines with electrons during the radiation dominante era (**references** [3]). Without this barrier there will be no suppression of inelastic reactions, in which O^{--} binds with nuclei. After recombination the OBe gas will undergo a decoupling with the plasma and the radiation, after that there is an adiabatic damping separation which remove the density fluctuation at smaller scale than the horizon of events. **This is the Warmer than cold dark matter model.**  know there is some OBe in the stars, but the thermonuclear **process** in the stars **interacts** with it and create anomalous isotopes. OBe can be ionized, but the more part of it is neutral. In the atmosphere, OBe atoms are the most part of the dark matter, in the ionosphere the particules are ionized by the radiation and neutralized by electron capture.

5 Conclusion

The O-Helium, this **particule** composed by helium nuclei and a doubled charged lepton O^{--} , is good hypothesis for explain the nature of the dark matter, and **it existence has been desmonstrated in the LHC, and the physicals characteristics too.** 

The warmer than cold dark matter scenario has the advantage to have a few parameters for explain the Universe, and the scenario is not very different than the cold dark matter scenario. The OHe dark matter can explain the observations from the center of the galaxies.

The problem is this model run successfully only if there is a strong Coulomb barrier in

the OHe particule. To confirm it existence, this needs a new research in atomics physics, astrophysics and simulation.

6 References

References

- [1] J-R.Cudell, M.Yu Khlopov, and Q. Wallemacq. Effects of dark atoms excitations. *Mod.Phys.Lett*, 2014.
- [2] M.Yu Khlopov. 10 years of dark atoms of composite dark matter. *Mod.Phys.Lett*, 2015.
- [3] J-R.Cudell, M.Yu Khlopov, and Q. Wallemacq. Some potential problems of ohe composite dark matter. *Bled Workshops in Physics 15*, 2014.

