

Problems of OHe dark matter

Steve Branchu

7 August 2017

Contents

1	Introduction	1
2	The origin of the model	2
2.1	Some particles dark matter candidates	2
2.2	The composite dark matter	2
3	The atomic structure of OHe	3
4	The OHe model of the Universe	5
5	Signatures of OHe	9
5.1	The solution for puzzles of direct dark matter searches	9
5.2	Positron annihilation in the galactic bulge	10
5.3	Composite dark matter solution for high energy positron excess	11
5.4	The LHC probes	11
6	Some potential problems	12
6.1	Inelastic process	12
6.2	Problems of OBe dark matter	15
6.3	Problem of seasonal variation for OHe model	16
7	Conclusion	16
8	References	17

1 Introduction

The dark matter, which corresponds to 25 % of total cosmological density, makes it possible to understand the creation of the large scale structures in the Big Bang. This matter is named like this because it is nonluminous. The existence of dark matter was confirmed by the astronomical observations. There are more accurate methods to calculate the mass of matter in the Universe. Two of them permit to discover the dark matter for the first time : the speed of rotations of the galaxies, which come from the Newton's laws, and the luminosity of the galaxies, which relates the mass of the galaxy to its luminosity. The data obtained by this 2 methods have not the same value that means there is missing matter, i.e dark matter. The non baryonic nature of the dark matter comes from analysis of the nucleosynthesis of the Big Bang and the cosmic microwave background anisotropies. The results obtained indicate the dark matter should be stable, saturate matter density and decouple from the plasma just before matter dominance era ([1]).

Today we don't know the nature of the dark matter, i.e its physical composition, the mass of its constituents and the interaction with other particles. For many models of dark matter, simulations of the large scale structure (LSS) formation have been done, and the results give different evolution of the LSS according for the model of the dark matter. There are several types of particles which can be the dark matter ([1]) : the simplest variant of dark matter is the Weakly Interactive Massives Particles (WIMP), the neutrino dark matter (which corresponding to the Hot dark matter scenario), the axions, the self-interacting dark matter, the decaying dark matter, the Teraparticles, the Walking technicolor models, the AC-leptons and the stable particles of 4th generation of quarks and leptons (discussed in section 2). The study of physical composition of dark matter is done by various direct, indirect methods and the accelaretors. The gravitationnal lensing which has the advantage to not depend on the frequency of the light, and the background of the galaxies not being punctual, the deflection varies across the source. Then there is dark matter captured in the Sun and the Earth, which emits neutrino which can be detected on the Earth, and finally the annihilation and decay of dark matter. The data obtained can be use for the searches for some dark matter particles in the LHC.

It is presented here one of the propositions for the cosmological dark matter, which is the simplest model of dark atoms, the O-Helium dark matter, abbreviated in OHe, which corresponds to the composite dark matter model. I will explain in this referat the research about OHe, the advantages of this model and finally the problems of this model.

2 The origin of the model

2.1 Some particles dark matter candidates

The axions, a wide class of particle models which possesses a symmetry breaking pattern, which can be described by pseudo-Nambu–Goldstone (PNG) field ([2]). The coherent oscillations of this field represent a specific type of CDM in spite of a very small mass of PNG particles $m_a = \Lambda^2/f$, where $f \gg \Lambda$, since these particles are created in Bose-Einstein condensate in the ground state, i.e. they are initially created as nonrelativistic in the very early Universe. This feature can be the general feature for all the axion-like PNG particles.

The self interacting dark matter, composed by mirror particles which have equivalence of left- and right-handed co-ordinate systems in the presence of P- and C- violation in weak interactions, should be strictly symmetric by their properties to their ordinary twins. After discovery of CP-violation it was shown by I. Yu. Kobzarev, L. B. Okun and I. Ya. Pomeranchuk in [3] that mirror partners cannot be associated with antiparticles and should represent a new set of symmetric partners for ordinary quarks and leptons with their own strong, electromagnetic and weak mirror interactions. It means that there should exist mirror quarks, bound in mirror nucleons by mirror QCD forces and mirror atoms, in which mirror nuclei are bound with mirror electrons by mirror electromagnetic interaction.

The decaying dark matter, composed by decaying particles with lifetime τ , exceeding the age of the Universe t_U , can be treated as stable. The primordial stable particles survives to the present time should be present in the modern Universe. However, even small effect of their decay can lead to significant contribution to cosmic rays and gamma background ([4]). The Unstable Dark matter scenarios [5] implied weakly interacting particles that form the structure on the matter dominated stage and then decay to invisible modes after the structure is formed.

The stable particles of 4th generation of quarks and leptons, which follows from heterotic string phenomenology and its difference from the three known light generations can be explained by a new conserved charge, possessed only by its quarks and leptons. Strict conservation of this charge makes the lightest particle of 4th family (neutrino) absolutely stable, but it was shown in [6, 7, 8] that this neutrino cannot be the dominant form of the dark matter.

2.2 The composite dark matter

The particles in the composite dark matter model can be electrically charged, but they are hidden in atom-like states maintaining dark matter of the modern Universe. This is the origin of the name of this model : the electrically charged constituents of dark atoms

may be not only elementary particles, but can be composite objects.

This idea was proposed by Sheldon Glashow in his model ([9]) : it is based on million times heavier partners of normal quarks and leptons related by a strict symmetry. The lightest of these partners are the tera-electrons and tera-U-quark, they could form a stable tera-helium atom (UUU)EE, in which +2 charged quark cluster (UUU) was bound by ordinary Coulomb force with two tera-electrons. It was proposed that in the early Universe, the excessive U-quarks first bind in (UUU) cluster, which recombines then with excessive tera-electrons to form tera-helium atom. As revealed in [10], there is an unrecoverable problem : binding of U-quarks and tera-electrons is incomplete, that implies an overproduction +1 and +2 charged like (Uud), (UUu) hadrons or (UUU)E ions, which bind with ordinary electrons. It makes impossible to realize the dark atom scenario not only in Glashow's sinister model, but also in any other model predicting stable +1 and -1 charged species.

Since 2006, different solutions for dark atom scenario were proposed ([11, 12, 13, 14, 15]), in which the important role of stable -2 charged species was revealed. The OHe dark atoms scenario was first proposed in 2005 ([11]) : in this model, dark atoms are composed by stable -2 charge particles, which should be heavy ($\sim 1TeV$) called O^{--} , which are bound by the Coulomb interaction with primordial helium. The candidates for O^{--} are : (a) clusters of heavy quarks formed by 3 stable anti-U quarks of fourth generation ([16, 7, 17]), (b) AC-leptons predicted in the extension of standard model ([18]) in the case of the almost-commutative geometry, (c) Technileptons and technibaryons in the framework of walking technicolor models (WTC) ([19]), and finally (d) the charged clusters, composed by 3 stable anti-quarks of fifth family ([3]).

The creation of OHe dark matter follows the scenario below : after its formation in the Standard Big Bang Nucleosynthesis (SBBN), 4He screens the excessive O^{--} charged particles in composite O-Helium. In all the considered forms of OHe, O^{--} behaves either as a lepton or as a heavy quark cluster with strongly suppressed hadronic interaction. The main features of OHe scenario are determined by the interaction with matter of helium shell of OHe. Existence of barrier provides dominance of elastic collisions, which is necessary for a self-consistent scenario.

3 The atomic structure of OHe

An OHe atom is composed by a helium nucleus and a heavy double charged particle O^{--} ([20]), which is a particle with strongly suppressed hadronic interaction. There are the mass M_O and M_{He} , and the charge $Z_O = 2e$. The potential interaction follows the laws below :

$$V(r \geq R_{He}) = -\frac{Z_O Z_{He} \alpha}{r} \quad (1)$$

$$V(r \leq R_{He}) = -\frac{Z_O Z_{He} \alpha}{r} \left(3 - \frac{r^2}{R_{He}^2} \right) \quad (2)$$




Figure 1 – The O-helium interaction potential (red) and the elementary Coulomb potential (green) obtained for a point-like helium nucleus, as a function of the distance r between O^{--} and the center of the helium nucleus.

where $\alpha = e^2/4\pi$ is the fine structure constant and r the distance between O^{--} and the center of the nucleus. The potential is represented in Figure 1. Then, we put the expression of potential (1) in the radial time-independent Schrödinger equation, to obtain the eigenvalues of an OHe atom. At angular momentum $l = 0$, the non-modified WKB approximation was used in ([21]) as potential V is regular at the origin ($\lim_{r \rightarrow 0} rV(r) = 0$). For states at $l \neq 0$, we replace $l(l+1)$ by $(l + \frac{1}{2})^2$ in the centrifugal term of the effective potential. For the first ten values of the quantum numbers n and l , the formula looks like pure hydrogen :

$$E_n^H = -\frac{1}{2}M_{He} \frac{(Z_O Z_{He} \alpha)^2}{n^2}$$

For the fundamental level, $E_1^H = 1.5879 MeV$. In fact the potential is no longer $\propto 1/r$, so energy depends on both n and l , the energy of ground state increases with E_1^H , when we take into account the charge distribution of the helium nucleus. The eigenvalues $E_{n,l}$ are shown in Table 1. In general, at fixed n , the pure hydrogen-like energy levels constitute a lower limit to which the levels $E_{n,l}$ tend as l increases, i.e. as the states are excited and thus as O^{--} and He lie further apart from each other, making the helium nucleus increasingly point-like. As it is well known that the WKB approximation is less accurate

for the deeper bound states, the energy of the ground state was computed by a variational method using up to 11 hydrogen-like s-orbitals and found the result -1.1771 MeV, which shows that the error on the values given in Table 1 is less than 0.1% ([20]).

n	$l = 0$	$l = 1$	$l = 2$	$l = 3$	$l = 4$	E_n^H
1	-1.1760	-	-	-	-	-1.5879
2	-0.3446	-0.3969	-	-	-	-0.3970
3	-0.1607	-0.1764	-0.1764	-	-	-0.1764
4	-9.2538^{-2}	-9.9230^{-2}	-9.9240^{-2}	-9.9239^{-2}	-	-9.9244^{-2}
5	-6.0057^{-2}	-6.3511^{-2}	-6.3511^{-2}	-6.3511^{-2}	-6.3510^{-2}	-6.3516^{-2}
6	-4.2097^{-2}	-4.4106^{-2}	-4.4106^{-2}	-4.4106^{-2}	-4.4106^{-2}	-4.4108^{-2}
7	-3.1136^{-2}	-3.2404^{-2}	-3.2404^{-2}	-3.2404^{-2}	-3.2404^{-2}	-3.2406^{-2}
8	-2.3957^{-2}	-2.4808^{-2}	-2.4811^{-2}	-2.4810^{-2}	-2.4810^{-2}	-2.4811^{-2}
9	-1.9002^{-2}	-1.9602^{-2}	-1.9602^{-2}	-1.9602^{-2}	-1.9602^{-2}	-1.9604^{-2}
10	-1.5439^{-2}	-1.5878^{-2}	-1.5878^{-2}	-1.5878^{-2}	-1.5878^{-2}	-1.5879^{-2}
n	$l = 5$	$l = 6$	$l = 7$	$l = 8$	$l = 9$	E_n^H
1	-	-	-	-	-	-1.5879
2	-	-	-	-	-	-0.3970
3	-	-	-	-	-	-0.1764
4	-	-	-	-	-	-9.9244^{-2}
5	-	-	-	-	-	-6.3516^{-2}
6	-4.4105^{-2}	-	-	-	-	-4.4108^{-2}
7	-3.2403^{-2}	-3.2406^{-2}	-	-	-	-3.2406^{-2}
8	-2.4810^{-2}	-2.4810^{-2}	-2.4810^{-2}	-	-	-2.4811^{-2}
9	-1.9604^{-2}	-1.9602^{-2}	-1.9602^{-2}	-1.9602^{-2}	-	-1.9604^{-2}
10	-1.5878^{-2}	-1.5878^{-2}	-1.5878^{-2}	-1.5878^{-2}	-1.5879^{-2}	-1.5879^{-2}

Table 1 – Energy levels $E_{n,l}$ (MeV) of the OHe atom, for the first ten values of the principal quantum number n and the corresponding angular momenta $l = 0, \dots, n - 1$. In the last column are also shown the pure hydrogen-like solutions E_n^H (MeV) obtained when the helium nucleus is assumed to be point-like. The exponents indicate the power of 10 by which the numbers have to be multiplied to obtain the energy in MeV ([20]).

4 The OHe model of the Universe

During the radiation dominance (RD) era, O-Helium, plasma and radiation are in thermal equilibrium, while the plasma transfers the energy and momentum to OHe ([1]). The radiation pressure in the plasma is transferred to density fluctuations to the OHe gas and transforms them in acoustic waves at scales up to the size of the horizon. To explain the

next step of the scenario, we use the following relations :

$$v = \sqrt{\frac{2T}{m_p}} \quad (3)$$

$$\sigma \approx \sigma_0 \sim \pi R_0^2 \approx 10^{-25} \text{cm}^2 \quad (4)$$

The formula (3) is the baryon thermal velocity, with T the temperature of the plasma and m_p the mass of the proton and the relation (4) is the cross section of OHe. At temperature $T \leq T_{od} \approx 200 S_3^{2/3} \text{eV}$ the energy and momentum transfer from baryons to O-Helium is not effective because :

$$n_B \langle \sigma v \rangle (m_p/m_0) t \leq 1$$

where $S_3 = m_0/(1 \text{TeV})$, and m_0 the mass of the OHe atom. In this formula $\sigma(m_p/m_0)$ is the transport cross section per nucleon. This relation means at this temperature the baryon thermal velocity is too weak and the transport cross section per nucleon is too small to make possible the transfer of energy to OHe atoms ([11]). After OHe gas decoupled from plasma, it started to become dominant after $t \sim 10^{12} \text{s}$ at $T \approx 1 \text{eV}$ and OHe atoms have the main role in the development of gravitationnal perturbations, and the formation of large scale structure. The nature of OHe determines the features of the composite dark matter scenario. At $T \geq T_{RM}$, the density of OHe gas is $\rho_d = (T_{RM}/T) \rho_{tot}$. After we obtain the total mass of OHe gas within the cosmological horizon $t = l_h$:

$$M = \frac{4\pi}{3} \rho_d t^3 = \frac{4\pi}{3} \frac{T_{RM}}{T} m_{Pl} \left(\frac{m_{Pl}}{T} \right)^2$$

During the period of decoupling $T = T_{od}$ this mass depends strongly on the OHe mass S_3 and is given by :

$$M_{od} = \frac{T_{RM}}{T_{od}} m_{Pl} \left(\frac{m_{Pl}}{T_{od}} \right)^2 \approx 2 \times 10^{44} S_3^{-2} g$$

At $T = T_0$ the OHe forms and within the cosmological horizon the total mass is $M_0 = M_{od} (T_{od}/T_0)^3 = 10^{37} g$.

During the radiation dominance era, just before the decoupling, the propagation of the sound waves in the plasma is limited the Jeans length λ_J of the OHe gas. The relativistic equation of state is $p = \epsilon/3$, at the cosmological horizon the Jeans length is at the same order $\lambda_J = l_h/\sqrt{3} = t/\sqrt{3}$. After decoupling it decreases to $\lambda_J \sim v_0 t$, with $v_0 = \sqrt{2T_{od}/m_0}$, and the Jeans mass decreases to :

$$M_J \sim v_0^3 M_{od} \sim 3 \times 10^{-14} M_{od}$$

we expect, at scales $M \leq M_0$ a strong suppression of fluctuations, and an adiabatic damping of sound waves in the plasma at the radiation dominance era for scales $M_0 \leq M \leq M_{od}$. This suppression has not the same effect as the free streaming suppression in the Warm Dark Matter model, and its impact on the structure formation needs a special study in detailed numerical simulations. In WDM, the free-streaming scale corresponds to the mode that enters the horizon at the time when WDM particles become non-relativistic ([22]). In the case of O-Helium, its composite nature makes the large scale structure formation more close to CDM, but a bit Warmer. Due to the slight suppression of small scale fluctuations, the OHe model is called the Warmer than Cold Dark Matter model (WtCDM).

After the decoupling from the baryonic matter, the OHe gas does not follow the formation of baryonic astrophysical objects (stars, planets, molecular clouds...) and forms the dark matter halos of galaxies, the problem is OHe has too small interaction to follow baryonic matter, but sufficiently large interaction to make baryonic objects opaque for it.

To simulate the power spectrum of the WtCDM, I use the CLASS code developed by Julien Lesgourgues ([23]). This code was created for the simulation of more complicated cosmological scenario. The framework on the code is the following : I had in all modules of the source code the parameter for choose the OHe dark matter scenario in the input file. Then in the module "perturbations.c", which calculate the perturbations between baryons and photons ([24]), I had the same equations adapted for the perturbations between baryons and OHe :

Synchronous gauge

$$\dot{\delta}_{ohe} = -\theta_{ohe} - \frac{1}{2}\dot{h} \quad (5)$$

$$\dot{\theta}_{ohe} = -\frac{\dot{a}}{a}\theta_{ohe} + c_s^2 k^2 \delta_{ohe} + \frac{4\bar{\rho}_b}{3\bar{\rho}_{ohe}} an_e \sigma_T (\theta_b - \theta_{ohe}) \quad (6)$$

The variable δ is the density fluctuation, θ the divergence of fluid velocity, h the synchronous metric perturbations, ϕ and ψ the conformal Newtonian metric perturbations, a is the scale factor, k is the wavenumber of Fourier mode, $\bar{\rho}$ the density, n_e the electron number density, $\sigma_T = 0.6652 \times 10^{-24} cm^2$ the Thomson scattering cross section (given by $d\sigma/d\Omega = 3\sigma_T(1 + \cos^2(\theta))/16\pi$) and P_i is the conjugate momentum to comoving position x^i ($P_i = a(\delta_{ij} + \frac{1}{2}h_{ij})p^j$ in synchronous gauge and $P_i = a(1 - \phi)p_i$ in conformal Newtonian gauge, p_i is the proper momentum). Finally the subscript b and ohe correspond to the baryons and OHe.

The momentum transfer from the baryon component is represented by $an_e \sigma_T (\theta_b - \theta_{ohe})$ in the equations (6) and (8). Momentum conservation in Thomson scattering then implies that a term $(4\bar{\rho}_b/3\bar{\rho}_{ohe})an_e \sigma_T (\theta_b - \theta_{ohe})$ has to be added to the equation for $\dot{\theta}_{ohe}$.

Conformal Newtonian gauge

$$\dot{\delta}_{ohe} = -\theta_{ohe} - 3\dot{\phi} \quad (7)$$

$$\dot{\theta}_{ohe} = -\frac{\dot{a}}{a}\theta_{ohe} + c_s^2 k^2 \delta_{ohe} + \frac{4\bar{\rho}_b}{3\bar{\rho}_{ohe}} a n_e \sigma_T (\theta_b - \theta_{ohe}) + k^2 \psi \quad (8)$$

The square of the baryon sound speed is evaluated from :

$$c_s^2 = \frac{\dot{P}_{ohe}}{\dot{\rho}_{ohe}} = \left(1 - \frac{1}{3} \frac{d \ln T_b}{d \ln a}\right) \quad (9)$$

where μ is the mean molecular weight. The OHe temperature evolves according to :

$$\dot{T}_{ohe} = -2\frac{\dot{a}}{a}T_{ohe} + \frac{8}{3}\frac{\mu}{m_e}\frac{\bar{\rho}_b}{\bar{\rho}_{ohe}} a n_e \sigma_T (T_b - T_{ohe}) \quad (10)$$

Equation (10) follows from the first law of thermodynamics, $dQ = (3/2)d(P_{ohe})/\rho_{ohe} + P_{ohe}d(1/\rho_{ohe})$, with specific heating rate $\dot{Q} = 4(\bar{\rho}_b/\bar{\rho}_{ohe})a n_e \sigma_T k_B (T_b - T_{ohe})$.




Figure 2 – The power spectrum of Cold Dark Matter model (purple), the Warm Dark Matter model (green) and the Warmer than Cold Dark Matter model (blue).

The synchronous gauge corresponds at a coordinate system where only the spatial coordinates are pertubated and the comsic time corresponds at the proper time of a comoving observer. This gauge is used for the numerical calculations. In the newtonian gauge the scalar portion of the perturbed metric is diagonal, but the vector part is not

diagonal. In this gauge, the expansion is "saw" like isotropic. The area of constant time can be perturbed and the potential is used in the equation of Poisson, i.e the potential assimilable at the newtonian potential at small scale.


The result is presented on the figure 2. The power spectrum of the warmer than cold dark model is as we expected, i.e between the power spectrum of warm dark matter and cold dark matter.

5 Signatures of OHe

5.1 The solution for puzzles of direct dark matter searches

Dark atom interpretation of the puzzles of direct dark matter search is based on the interactions between OHe nucleus and the baryonic matter. If dark matter can bind to normal matter, the results of DAMA/NaI and DAMA/LIBRA experiments ([25]) could be interpreted like come from the radiative capture of thermalized OHe and could depend on the detector composition and temperature ([26]). The concentration of OHe in the underground detector is determined by the equilibrium between the infalling cosmic OHe flux and its diffusion to the center of Earth. The infalling flux experiences annual changes due to Earth's rotation around Sun and modifies the OHe concentration, so this local OHe concentration possess annual modulations.

Many experiments have been done but only one gave a positive result ([26]) : DAMA/NaI and DAMA/LIBRA. The interpretation of the experiments in terms of OHe model are based on the idea that OHe slowed down in the terrestrial matter, can form a few keV bound state with sodium nucleus, if OHe is situated beyond the nucleus. The positive result of these experiments is explained by annual modulation in reaction of radiative capture in the detector :



The low energy OHe-nucleus bound states is possible only for intermediate-mass nuclei : this explain the negative results of LUX experiments and all heavy nuclei detectors (XENON 100, XENON 1T etc...). To calculate the rate of this capture, [27] use an analogy with the radiative capture of neutron by proton, without the M1 transition from conservation of orbital momentum and suppression of E1 transition in the case of OHe. Since OHe is isoscalar, isovector E1 transition can take place in OHe-nucleus system only due to effect of isospin nonconservation, which can be measured by the factor f (explained after equation (11)). The rate of OHe radiative capture according to atomic numbers A and Z, the energy level E in the medium and the temperature T is given by the relation

below :

$$\sigma v = \frac{f\pi\alpha 3T}{m_p^2 \sqrt{2Am_p E}} \left(\frac{Z}{A}\right)^2 \quad (11)$$

with the factor $f = (m_n - m_p)/m_N \approx 1.4 \times 10^{-3}$, corresponding to the difference of mass of neutron m_n and proton m_p , relative to the mass of nucleon m_N . The capture rate is proportional to the temperature at the thermal equilibrium : this leads to a suppression of this effect in cryogenic detectors. Since OHe capture rate is proportional to the temperature, it looks like it is suppressed in cryogenic detectors by a factor of order 10^{-4} . However, for the size of cryogenic devices less, than few tens meters, OHe gas in them has the thermal velocity of the surrounding matter and the suppression relative to room temperature is only $\sim m_A/m_O$ (m_O is the mass of OHe). Then the rate of OHe radiative capture in cryogenic detectors is given by equation (11), in which room temperature T is multiplied by factor m_A/m_O , and the equation becomes :

$$\sigma v = \frac{f\pi\alpha 3T}{m_p^2 \sqrt{2Am_p E}} \left(\frac{Z}{A}\right)^2 \times \frac{m_A}{m_O} \quad (12)$$

5.2 Positron annihilation in the galactic bulge

The composite nature of O-helium dark matter results in a number of observable effects. The satellite INTEGRAL observed a positron-annihilation line excess in the galactic bulge. This observation can be explained by the presence of OHe ([20, 26]). The O-helium collisions excite the 2S level, after OHe de-excites by electron-positron pair-production. The rate of collisions between OHe particles can take place everywhere in the Galaxy. It is nonzero and grows in the regions of higher OHe density, particularly in the central part of the Galaxy because these collisions are enhanced. The collision rate is estimated to ([28]) :

$$\frac{dN}{dt} = n^2 \sigma v_h \frac{4\pi}{3} r_b^3 \approx 3 \times 10^{42} S_3^{-2} s^{-1}$$

At a velocity $v_h \sim 3 \times 10^7 cm/s$, the energy transfered in the collisions is $\Delta E \sim 1 MeV S_3$. The collisions excite the OHe gas, which de-excites by pair production, then the de-excitation in E0 transitions had been observed by excess of positron-annihilation line. However, measurement of the collisions rate depends of the density of OHe, but the theoretical estimations for dark matter are uncertain because in the bulge OHe gas density doesn't lead to strong gravitationnal effect and can vary by several orders of magnitude. The last analysis indicates a lower value of dark matter density, which makes it possible to explain the observed effect if the mass of O^{--} near 1.0 TeV.

The collisions excite the OHe gas at the first level with non-zero angular momentum,

the E1 transition create gamma lines with principal quantum numbers n and m , when n is higher than m , at the energies :

$$E_{nm} = \left(\frac{1}{m^2} - \frac{1}{n^2} \right) \times 1.5879 \text{ MeV}$$

The interest of this predictions is to analyze and compare the unidentified lines from the center of the Milky Way. In all line emissions from the Galaxy ([20]), there are some X-Ray and gamma ray, probably produced by OHe. The important line for INTEGRAL data have an energy around 20 keV, and the X-Ray lines have a lower energy. XMM-Newton can check the observations between 0.1 – 12keV.

Taking into account all the possible electric dipole transitions (E1) between the states of OHe between 3 and 4 keV, several hundreds of allowed lines were found in ([20]), with energies from the eV to the MeV range. The comparison between the predictions and the observations provides an effective tool to test OHe dark matter model.

5.3 Composite dark matter solution for high energy positron excess

PAMELA and AMS02 experiments ([29, 30, 31]) found an excess of high-energy cosmic-ray positrons. This can be explained by the decay of doubly positive charged techniparticles to pairs of same-sign leptons. This is the two-component dark atom model, based on Walking Technicolor sparse WIMP-like component of atom-like state, made of positive and negative doubly charged techniparticles. This is possible for the mass of decaying +2 charged particle below 1 TeV and depends on the branching ratios of leptonic channels ([26]). Since even pure lepton decay channels are inevitably accompanied by gamma radiation the important constraint on this model follows from the measurement of cosmic gamma ray background in FERMI/LAT experiment. The multi-parameter analysis of decaying dark atom constituent model is under way in order to determine the maximal model independent value of the mass of decaying +2 charge particle, at which this explanation is possible.

5.4 The LHC probes

One of the propositions to understand the constitution OHe atom, is to do searches for stable doubly charged lepton-like particles at the LHC ([26]). The objective is to compare the results with astroparticle data. The ATLAS and CMS experiments give the lower value for double charged particle around 700 GeV ([32]), this result will permit to test OHe explanation when the future data can approach 1 TeV range in this searches. Recently, a Yu S Smirnov's studies improved this result, he found a value for double charged particle around 685 GeV ([33]).

6 Some potential problems

6.1 Inelastic process

It was first assumed that the effective potential between OHe and a normal nucleus would have a barrier, preventing He and/or O^{--} from falling into the nucleus, allowing only one bound state, and decreases rapidly the interactions of OHe. Under these conditions elastic collisions dominate in OHe interactions with matter, and lead to a successful OHe scenario. The cosmological and astrophysical effects of such composite dark matter (dark atoms of OHe) are dominantly related to the helium shell of OHe and involve only one parameter of new physics, the mass of O^{--} . In this section, I want to explore another scenario ([11, 14]), in which OHe dark matter interacts strongly with normal matter : OHe is neutral, but a priori it has an unshielded nuclear attraction to matter nuclei. I explain in this section the consequences of this effect.

At the beginning of the Universe, inelastic scattering between particles of OHe and between OHe and primordial He decreased the quantity of OHe ([34]). The nuclear reactions are :



where Be is the beryllium. The OHe forms at a temperature T_0 which depends of its binding energy, which is 1.175 MeV, that corresponds to $T_0 = 50\text{keV}$, i.e the energy absorbed during the formation of OHe. The cosmological time is calculated with the temperature, the inelastic process start at a time :

$$t(s) \simeq \frac{1}{T^2(MeV)} \simeq \frac{1}{0.05^2} = 400s$$

after the Big Bang and continue until helium freezes out at $t_* \simeq 600s$.

During these 200 s, the rate of the quantity of OHe decreases like this :

$$\frac{dn_{OHe}}{dt} = -3Hn_{OHe} - n_{OHe}^2\sigma_1v_1 - n_{OHe}n_{He}\sigma_2v_2 \quad (15)$$

with n_{OHe} and n_{He} are the number densities of OHe and He, $H = 1/2t$ the expansion rate of the Universe during the radiation dominance era, σ_1 , σ_2 , v_1 and v_2 are respectively the cross section and the relative velocity of the reaction (13) and (14). The only factor which affect n_{He} is the expansion :

$$\frac{dn_{He}}{dt} = -3Hn_{He} \iff n_{He}(t) = n_{He}^0 \left(\frac{t_0}{t} \right)^{3/2} \quad (16)$$

where n_{He}^0 is the number density of He at $t = t_0$. We calculate the fraction of free OHe atoms due to their inelastic reactions, with the ratio f of the number density of OHe and He nuclei, $f = \frac{n_{OHe}}{n_{He}}$. The expressions (15) and (16) give the relation of evolution :

$$\frac{df}{dt} = -n_{He}f(\sigma_1 v_1 f + \sigma_2 v_2) \quad (17)$$

The cross sections σ_1 and σ_2 depend of the size of the nucleus :

$$\sigma_1 \approx 4\pi(2r_{OHe})^2 \quad (18)$$

$$\sigma_2 \approx 4\pi(r_{OHe} + r_{He})^2 \quad (19)$$

where r are the radius of respectively the OHe nucleus and the He nucleus. The both are approximately equal to 2 fm, $\sigma_1 \approx \sigma_2 \approx 64\pi 10^{-26} cm^2$. The mean relative velocities v_1 and v_2 are obtained by the Maxwell-Boltzmann velocity distributions, because the species are in thermal equilibrium with the plasma :

$$v_1 = \sqrt{\frac{8T}{\pi\mu_1}} \quad (20)$$

$$v_2 = \sqrt{\frac{8T}{\pi\mu_2}} \quad (21)$$

with $\mu_1 = \frac{m_{OHe}}{2}$ the reduced mass of OHe-OHe and $\mu_2 \simeq m_{He}$ the reduced mass of OHe-He. During the radiation-dominated era, the relation between time and temperature is : $Tt^{1/2} = T_0 t_0^{1/2}$, we insert it in the expression of velocities, and use the relation (17) to obtain :

$$\frac{df}{dt} = -\gamma \frac{1}{t^{7/4}} f(\alpha f + \beta) \quad (22)$$

with

$$\alpha = \frac{\sigma_1}{\sqrt{\mu_1}} \quad (23)$$

$$\beta = \frac{\sigma_2}{\sqrt{\mu_2}} \quad (24)$$

$$\gamma = n_{He}^0 t_0^{7/4} \sqrt{\frac{8T_0}{\pi}} \quad (25)$$

The initial condition is $f(t_0) = f_0$, the solution of (22) is :

$$f(t) = \frac{\beta f_0}{\exp(\frac{4}{3}\beta\gamma(t_0^{-3/4} - t^{-3/4}))(\alpha f_0 + \beta) - \alpha} \quad (26)$$

We can calculate the number density of He at the beginning of the baryon nucleosynthesis from its actual value. It corresponds today about 10% of all baryons, and the current critical density :

$$\rho_c^1 = 5.67 \times 10^{-6} m_p / cm^3 \iff n_{He}^1 \simeq 2.8 \times 10^{-8} cm^{-3} \quad (27)$$

Then, it is necessary to add the effect of the expansion, which is proportionnal at the temperature of the CMB $n_{He}^0 = n_{He}^1 \left(\frac{T_0}{T_1} \right)^3 \simeq 2.8 \times 10^{-17} cm^{-3}$. At the early Universe the number of O^{--} particles was equal to OHe particles. Between t_0 and today, O^{--} particles may have been bound in different structures, but they have not been created or destroyed, so that their number density has only been diluted by the expansion in the same way as that of He nuclei, so that the ratio of the number density of O^{--} particles to the number density of He nuclei remains unchanged : $\frac{n_O^0}{n_{He}^0} = \frac{n_O^1}{n_{He}^1}$. The fraction of OHe hasn't been change, almost 25% of the critical density, so we can calculate the parameter $f_0 \simeq 0.05$. Now it is possible to insert this value in the equation (26) and calculate the fraction at the freeze-out time :

$$f(t_*) \simeq 5 \times 10^{-6133} \ll f_0 \quad (28)$$

this results means the OBe creation by the reaction 1 decreases rapidly. To be more clearly, the reaction (14) is prevalent, so the dark matter is now in form of OBe. The suppression of f comes when the exponential term in (26) is evaluated to be e^{14127} . This value represents the number N_2 of reactions that happened between t_0 and t_* :

$$\begin{aligned} N_2 &= \int_{t_0}^{t_*} n_{He}(t) \sigma_2 v_2(t) dt \\ &= n_{He}^0 t_0^{3/2} \sigma_2 \sqrt{\frac{8T_0 t_0^{1/2}}{\pi \mu_2}} \int_{t_0}^{t_*} \frac{1}{t^{7/4}} dt \\ &= n_{He}^0 t_0^{7/4} \sqrt{\frac{8T_0}{\pi}} \frac{\sigma_2}{\sqrt{\mu_2}} \left(-\frac{4}{3} \right) \left(\frac{1}{t_*^{3/4}} - \frac{1}{t_0^{3/4}} \right) \\ &= \frac{4}{3} \beta \gamma \left(\frac{1}{t_0^{3/4}} - \frac{1}{t_*^{3/4}} \right) \end{aligned}$$

This relation means the realization of the OHe dark matter model implies a very strong suppression of the reaction (14), which corresponding to $N_2 \ll 1$. This particular case needs the development of a strong dipole Coulomb barrier in OHe-He interaction.

6.2 Problems of OBe dark matter

Due to the Coulomb barrier, OBe can't capture helium nucleus, and it recombines with electrons during the radiation dominante era ([34]) at the temperature $T_{od} = 2eV$. Without this barrier there will be no suppression of inelastic reactions, in which O^{--} binds with nuclei. It makes anomalous helium the dominant form of dark matter in this scenario. After recombination the OBe gas will undergo a decoupling from the plasma and the radiation, after that there is an adiabatic damping slightly suppresses density fluctuation at scales smaller than the scale of the horizon in the period of He recombination.

This dark matter is mixed with ordinary matter in the process of galaxy formation, and become collisional on the scale of the galaxies. So there is some OBe in the stars, but the thermonuclear reaction in the stars interacts with it and create anomalous isotopes. OBe can be ionized, but the dominant part is neutral. In the atmosphere, OBe atoms are the prevalent part of the dark matter, in the ionosphere the particles are ionized by the radiation and neutralized by electron capture. When OBe fall down on the Earth there are many collisions which decrease the velocity of OBe. The cross section of OBe is in the order of $\sigma \approx 10^{-15} - 10^{-14} cm^{-2}$, with the number density of terrestrial matter $n = 0.27 \times 10^{23} \text{molecules/cm}$ we calculate the OBe atoms velocity inside the Earth at :

$$V = \frac{g}{n\sigma v} \leq 2.7 \times 10^{-11} cm/s \approx 270 fm/s \quad (29)$$

To determine the OBe abundance in the Earth we need to know the equilibrium between the in-falling and down-drifting fluxes. The flux of OBe in-falling from dark matter halo is given by :

$$F = \frac{n_0}{8\pi} \times |\bar{V}_h + \bar{V}_E| \quad (30)$$

with $V_h = 220 km/s$ the speed of the Solar System, $V_E = 29.5 km/s$ the speed of the Earth and $n_0 = 3 \times 10^{-4} cm^{-3}$ is the assumed local density of OBe dark matter. To simplify the calculation, Cudell et al ([34]) didn't take into account the annual modulation of the incoming flux and take $|\bar{V}_h + \bar{V}_E| = u \approx 300 km/s$. The equilibrium concentration of OBe is obtained by :

$$n_{0E} = \frac{2\pi \times F}{V} \quad (31)$$

and the ratio of anomalous isotopes to the total amount of the Earth matter is given by :

$$r_{0E} = \frac{n_{0E}}{n} = \frac{2\pi \times F\sigma v}{g} \geq 3.1 \times 10^{-9} \quad (32)$$

The upper limits on the anomalous helium abundance are very stringent ([35]) $r_{0E} \leq 10^{-19}$, and the rough estimate is ten orders of magnitude too large. Together with the

other problems of OBe Universe stipulated above, this rules out the OBe scenario.

6.3 Problem of seasonal variation for OHe model

O-Helium fall down to the terrestrial surface with the same velocity as OBe (29). At a depth L below the Earth's surface, the drift timescale is $t_{dr} \sim L/V$ where V is given by equation (29). It means the motion of the Earth around the Sun cause a change of the incoming flux, should lead at the at the depth $L \sim 10^5\text{cm}$ ([14]) to the corresponding change in the equilibrium underground concentration of OHe on the timescale $t_{dr} \approx 2.5 \times 10^2 S_3^{-1}\text{s}$ (S_3 is defined in the section 4). In underground detectors, OHe dark atoms are slowed down to thermal energies and give rise to energy transfer $\sim 2.5 \times 10^{-4}\text{eV}A/S_3$, where $A \sim 30$ is the average atomic weight in terrestrial surface matter. OHe slowed down far below the threshold for direct dark matter detection. The equilibrium concentration of OHe is obtained by :

$$n_{0E} = \frac{2\pi \times F}{V} = n_{0E}^{(1)} + n_{0E}^{(2)} \times \sin(\omega(t - t_0)) \quad (33)$$

where $\omega = 2\pi/T$, $T = 1\text{year}$ and t_0 is the phase. This parameters are a consequences of the rotation of the Earth. The 2 other variables are the average concentration

$$n_{0E}^{(1)} = \frac{n_0}{320S_3A^{1/2}}V_h \quad (34)$$

and the annual modulation of concentration

$$n_{0E}^{(2)} = \frac{n_0}{640S_3A^{1/2}}V_E \quad (35)$$

The rate of nuclear reactions of OHe with nuclei is proportional to the local concentration and the energy release in these reactions should lead to observable signal. There are two parts of the signal : the one determined by the constant part and the second determined by annual modulation, which is concerned by the strategy of dark matter search in DAMA experiment ([36, 37, 38, 39, 40]).

The terrestrial matter is opaque for OHe, what should inevitably lead to an effect of Earth matter shadowing for the OHe flux and corresponding diurnal modulation. This effect needs special study in the confrontation with the constraints, recently obtained in DAMA/LIBRA experiment ([26, 25]).

7 Conclusion

The O-Helium dark atom, composed by helium nuclei and a double charged O^{--} particle, is a serious candidate for the dark matter. The physical nature of OHe is at the center of

the experimental astroparticles searches.

The warmer than cold dark matter scenario has the advantage to have a few parameters for explain the Universe, and the scenario is not very different from the cold dark matter scenario. The advantage is there is only one parameter in OHe scenario : the mass of OHe. The OHe dark matter can explain the observations from the center of the galaxies, especially the positron line excess due to OHe de-excitation. ATLAS experiments at LHC gave lower limits of the particles which compose OHe, but didn't rule it out yet.

The existence of heavy stable particles is one of the popular solutions for the dark matter problem. These particles are usually considered to be electrically neutral, but dark matter can also be formed by stable heavy charged particles bound in neutral atom-like states by the ordinary Coulomb attraction. The OHe dark matter model needs an in-depth study about OHe nuclear interaction and quantum mechanical proof for the existence of barrier.

8 References

References

- [1] M.Yu Khlopov. The puzzles of dark matter searches. *AIP Conf.Proc.*, 1241:388–397, 2009. arXiv:0911.5685.
- [2] S.G. Rubin M.Yu. Khlopov. *Cosmological pattern of microphysics in the inflationary universe*. Springer-Verlag New York Inc, 2004.
- [3] Anamarija Borštnik Bračič and Norma Susana Mankoč Borštnik. On the origin of families of fermions and their mass matrices. *Phys. Rev. Lett.*, 74:30, 2006. arXiv:hep-ph/0512062.
- [4] Christoph Weniger Alejandro Ibarra, David Tran. Indirect searches for decaying dark matter. *International Journal of Modern Physics A*, 28:47, 2013. arXiv:1307.6434.
- [5] M.Yu. Khlopov A.S. Sakharov. Horizontal unification as the phenomenology of the theory of 'everything'. *Phys.Atom.Nucl.*, 57:651–658, 1994.
- [6] K.I. Shibaev K.M. Belotsky, M.Yu. Khlopov. Sakharov's enhancement in the effect of 4th generation neutrino. *Grav.Cosmol.Suppl*, 6:140–143, 2000.
- [7] K.Shibaev K.Belotsky, M.Khlopov. Stable matter of 4th generation: hidden in the Universe and close to detection ? In *The 12th Lomonosov Conference on Elementary Particle Physics, Moscow, 25-31 August*, pages 180–184, 2005. arXiv:astro-ph/0602261.

- [8] M. Khlopov R.V. Konoplich K.M. Belotsky, D. Fargion. May Heavy neutrinos solve underground and cosmic ray puzzles ? *Phys.Atom.Nucl*, 71:147–161, 2008. arXiv:hep-ph/0411093.
- [9] S.L. Glashow. A sinister extension of the standard model to $SU(3) \times SU(2) \times SU(2) \times U(1)$. In *XI Workshop on Neutrino Telescopes, Venezia, February 22-25*, pages 539–547, 2005. arXiv:hep-ph/0504287.
- [10] D. Fargion and M. Khlopov. Tera-leptons shadows over sinister Universe. *Gravitation and Cosmology*, 19:219–231, 2013. arXiv:hep-ph/0507087.
- [11] M. Y. Khlopov. Composite dark matter from 4th generation. *JETP Lett.*, 83:1–4, 2006. arXiv:astro-ph/0511796.
- [12] Chris Kouvaris M. Y. Khlopov. Composite dark matter from a model with composite Higgs boson. *Phys.Rev.D*, 78:23, 2008. arXiv:0806.1191.
- [13] C. A. Stephan D. Fargion, M. Khlopov. Dark matter with invisible light from heavy double charged leptons of almost-commutative geometry? *Class.Quant.Grav.*, 23:7305–7354, 2006. arXiv:astro-ph/0511789.
- [14] Evgeny Yu. Soldatov Maxim Yu. Khlopov, Andrey G. Mayorov. The dark atoms of dark matter. *Prespacetime Journal*, 1:1403–1417, 2010. arXiv:1012.0934.
- [15] Maxim Yu. Khlopov. Physics of Dark Matter in the Light of Dark Atoms. *Modern Physics Letters A*, 26(38):2823–2839, 2011. arXiv:1111.2838.
- [16] K.M.Belotsky et al. Heavy hadrons of 4th family hidden in our Universe and close to detection ? *Gravitation and Cosmology*, 11:3–15, 2005. arXiv:hep-ph/0411271.
- [17] M.Yu.Khlopov. New symmetries in microphysics, new stable forms of matter around us. *Gravitation and Cosmology* 12, 1:39–49, 2006. arXiv:astro-ph/0607048.
- [18] Christoph A. Stephan. Almost-Commutative Geometries Beyond the Standard Model. *Journal of Physics A: Mathematical and Theoretical*, 39:9657–9670, 2006. arXiv:hep-th/0509213.
- [19] Dennis Dietrich, Francesco Sannino, and Kimmo Tuominen. Light composite Higgs and precision electroweak measurements on the Z resonance: An update. *Physical Review D*, 78:8, 2006. arXiv:hep-ph/0510217.
- [20] J-R.Cudell, M.Yu Khlopov, and Q. Wallemacq. Effects of dark atoms excitations. *Mod.Phys.Lett. A*, 29:8, 2014. arXiv:1411.1655.

- [21] N. Fröman and P.O. Fröman. *JWKB approximation : Contributions to the theory*. North-Holland Pub. Co, 1965.
- [22] Katarina Markovič and Matteo Viel. Lyman- α Forest and Cosmic Weak Lensing in a Warm Dark Matter Universe. *Publications of the Astronomical Society of Australia*, 31:20, 2013. arXiv:1311.5223.
- [23] Thomas Tram Diego Blas, Julien Lesgourgues. The Cosmic Linear Anisotropy Solving System (CLASS) II : Approximation schemes. *Journal of Cosmology and Astroparticle Physics*, 34:35, 2011. arXiv:1104.2933.
- [24] Edmund Bertschinger Chung-Pei Ma. Cosmological Perturbation Theory in the Synchronous and Conformal Newtonian Gauges. *Astrophysical Journal*, 455:45, 1995. arXiv:astro-ph/9506072.
- [25] R. Bernabei et al. [DAMA Collaboration]. DAMA/LIBRA-Phase1 results and Perspectives of the Phase2. In *Vth Roma International Conference on Astroparticle Physics, Noto, Sicily, Italy, 30 Sep-3 Oct*, volume 121, page 5, 2014.
- [26] M.Yu Khlopov. 10 years of dark atoms of composite dark matter. In *The XVIIIth Bled Workshop on Physics "What Comes Beyond the Standard Models ?" : Bled, Slovenia, July 11-19*, volume 16, pages 71–77, 2015. arXiv:1512.01081.
- [27] Maxim Yu Khlopov. Dark atoms and puzzles of dark matter searches. *International Journal of Modern Physics A*, 29:26, 2014. arXiv:1402.0181.
- [28] M. Y. Khlopov. Composite dark matter from stable charged constituents. In *Presented at 20th Rencontres internationales de Blois, Chateau de Blois, France*, 2008. arXiv:0806.3581.
- [29] C. Kouvaris K. Belotsky, M. Khlopov and M. Laletin. Decaying Dark Atom constituents and cosmic positron excess. *Advances in High Energy Physics*, 2014:20, 2014. arXiv:1403.1212.
- [30] M. Laletin K. Belotsky, M. Khlopov. Dark Atoms and their decaying constituents. In *Bled Workshops in Physics*, volume 15, pages 1–10, 2014. arXiv:1411.3657.
- [31] C. Kouvaris K. Belotsky, M. Khlopov and M. Laletin. High Energy Positrons and Gamma Radiation from Decaying Constituents of a two-component Dark Atom Model. *International Journal of Modern Physics D*, 24:11, 2015. arXiv: 1508.02881.
- [32] ATLAS Collaboration. Search for heavy long-lived multi-charged particles in pp collisions at 8 TeV using the ATLAS detector. *The European Physical Journal C*, 75:23, 2015. arXiv:1504.04188.

- [33] M.Yu Khlopov and Yu.S. Smirnov. Search for double charged particles as direct test for Dark Atom Constituents. In *VIA talk at Moscow astrophysical seminar*, 2017.
- [34] J-R.Cudell, M.Yu Khlopov, and Q. Wallemacq. Some potential problems of ohe composite dark matter. In *The XVIIth Bled Workshops in Physics on What Comes Beyond the Standard Models ? : Bled, Slovenia, July 20-28*, volume 15, pages 66–74, 2014. arXiv:1412.6030.
- [35] R. J. Holt Z.-T. Lu T. P. O'Connor J. P. Schiffer P. Mueller, L.-B. Wang. Search for anomalously heavy isotopes of helium in the Earth's atmosphere. *Phys. Rev. Lett.*, 92:13, 2004. arXiv:nucl-ex/0302025.
- [36] R. Bernabei et al. [DAMA Collaboration]. First results from DAMA/LIBRA and the combined results with DAMA/NaI. *The European Physical Journal C*, 56:333–355, 2008. arXiv:0804.2741.
- [37] R. Bernabei et al. [DAMA Collaboration]. New results from DAMA/LIBRA. *Eur. Phys. J. C*, 67:39–49, 2010. arXiv:1002.1028.
- [38] R. Bernabei et al. [DAMA Collaboration]. Particle Dark Matter in DAMA/LIBRA. In *Vulcano Workshop 2010 "Frontier Objects in Astrophysics and Particle Physics"*, *Vulcano (Italy)*, 2010. arXiv:1007.0595.
- [39] R. Cerulli et al [DAMA Collaboration]. DAMA annual modulation and mirror Dark Matter. *The European Physical Journal C*, 77:20, 2017. arXiv:1701.08590.
- [40] R. Bernabei et al [DAMA Collaboration]. DAMA/LIBRA results and perspectives. In *19th Bled Workshop "What Comes Beyond the Standard Models" : Bled, Slovenia, July 11-19*, volume 17, pages 1–7, 2016. arXiv:1612.01387.