

Report

Shadow matter with 4 generations of fermions

Nepochataya Olga

M17-115

Moscow, 2018

Content:

1. Introduction
2. Model
 - 2.1. Content of primary helium
 - 2.2. Baryosynthesis and leptosynthesis
3. Conclusion
4. Literature

1. Introduction.

Modern physics deals with three basic types of spatial symmetry: reflection, translation and rotation. For elementary particles only translation and rotation are performed, reflection (P - symmetry) is violated in a weak interaction. This fact was assumed theoretically by Lee and Yang, and then it was confirmed experimentally.

For example, in beta-decay only left-polarized neutrino are born, and reflection gives (P - transformation) neutrino, which can't take part in weak interaction.

P-violation means that left and right coordinate systems are not equivalent, and the Universe has preferred orientation of coordinate axes. Li and Yang assumed that there is a mirror partner for every elementary particle. In this case P-symmetry is not broken.

For a long time antiparticles were considered as candidates for the role of mirror particles. But the discovery of CP-violation showed that antiparticles are not mirror particles. I.Pomeranchuk, L.Kobzarev and L.Okun' demonstrated that mirror particles, if they exist, take part only in gravitational interaction.

2. Model.

In this paper, the possibility of existence mirror world is considered. The model suggests that this world exists parallelly with the usual world and consists of 4 generations of fermions (in the usual world there are 3 generations). We suggest that three generations of mirror particles are copies of three generations of usual particles, and the 4th generation consists of particles with masses less than mass of the “mirror Z-boson”, and 4th mirror generation lepton is lighter than corresponding neutrino.

We suggest that mirror world and usual world have different cosmology evolution. For example they had never been in the equilibrium. Shadow world interacts with the usual one only by gravity, but gravity is too weak for establish the thermodynamic equilibrium.

2.1. Content of primary helium.

The most important restriction for all models is connected with the data of content of primary helium (the ratio between quantity of ^4He and quantity of all nuclei in the end of the primary nucleosynthesis period (fig.1)). This data restrict the quantity of any relativistic matter (including shadow matter) on the moment $t \sim 1$ s, $T \sim 1$ MeV. The modern researches based on the cosmological nucleosynthesis and the cosmic microwave background gives the value $23,1\% < Y < 26,7\%$.




fig. 1. The ratio between quantity of chemical elements and quantity of all nuclei vs the quantity of barions. The thickness of lines corresponds to accuracy. Rectangles show the regions of values within the limits of errors from the experiments. Vertical stripes show regions of η_b values which are not forbidden by data of BBN and CMB. Index “p” means “primordial”.

With good accuracy the part of primordial helium formed in the early Universe is twice as many nucleons, which were neutrons at the moment of sufficient propagation of deuterium and the beginning of the process of formation of nuclei.

If there were 4th types of neutrino, the effective number of particle types would become 3.817 instead of 3.363, and the temperature drop rate increases $\sqrt{3.817/3.363} \approx 1.135$ -fold.

It increases the fraction of neutrons and helium.

Adding each new generation increases Y on the value ≈ 0.013 . It gives a restriction on the number of light neutrino generations. It can't be more than 4. Another restriction for the number of generations in usual world is connected with decay width of Z_0 -boson. The mass of the 4-th generation particles must be more than $\frac{1}{2}$ mass of Z_0 -boson, but this case is not considered in this paper.

Calculation of the amount of primary helium:

The ratio between amount of protons and neutrons:

$$\frac{n}{p} = \exp\left(-\frac{\Delta m}{T}\right)$$

$$\Delta m = M_n - M_p = 1.29 \text{ MeV}$$

Where M_p and M_n - masses of protons and neutrons, T - freezing temperature.

$$T \approx \frac{(kG)^{\frac{1}{6}}}{G_F^{\frac{2}{3}}} (3)$$

$$k = \sum g_s \left(\frac{T_i}{T}\right)^4 + \frac{7}{8} \sum \frac{g_s}{2} \left(\frac{T_i}{T}\right)^4,$$

k - statistical factor characterizing the density of the Universe. In our model we take into account usual and mirror photons, usual and mirror electrons and positrons, three generations of usual neutrino and four generations of mirror one. This factor for the Universe of our model is 6,88.

Temperature depends only on k . It means that the ratio between amount of protons and neutrons can be rewritten:

$$(n/p)' = (n/p)^{(k_0/k)^{\frac{1}{6}}}$$

k is the value we calculate for the usual world with three generations of fermions.

$$k_0 = 5,375$$

$$(n/p)' = 0,154$$

Then by chain of nuclear reactions most of neutrons became to the helium nuclei. The formation of heavier nuclei does not occur because of the Coulomb's barrier. Also there are no stable nuclei with masses 5 and 8.

Then the concentration of ${}^4\text{He}$ is

$$Y_p = \frac{He}{Barion} = \frac{2 \cdot (n/p)}{1 + n/p} \approx 0.267$$

This concentration falls within the constraints from the experimental data. It means that mirror world with 4 generations of fermions is possible.

2.2. Baryosynthesis and leptosynthesis.

The only conserved quantities in the $SU(3) \times SU(2) \times U(1)$ Standard Model correspond to gauge symmetries: electroweak generator T_3 , electroweak hypercharge Y , two generators of $SU(3)$ and B-L - the difference between baryonic and lepton charges (not this charges). In the period of thermodynamic equilibrium establishment, baryosynthesis and leptosynthesis, the ratio between baryonic quantity and B-L is calculated as follows:

$$N_b = N_{b-l} \cdot \frac{8 \cdot N_g + 4 \cdot N_d}{22 \cdot N_g + 13 \cdot N_d},$$

where N_g is a number of generations and N_d is the number of scalar doublets.

For three generations of usual matter this ratio is $28/79$, for three usual generations and four mirror generations - $16/45$. Adding new mirror generation doesn't change much this ratio. It means that mirror world with 4 generations of fermions is possible.

3. Conclusion.

In this paper the possibility of existing of mirror matter with 4 generations of fermions is described. It is shown that in the processes of baryosynthesis, leptosynthesis and nucleosynthesis there were no factors forbidding mirror matter with 4 generations of fermions.

The big part of warm dark matter can consist of mirror particles, because mirror particles take part only in gravitational interaction with usual matter. Shadow and usual matter have similar cosmological evolution and they can have similar total masses in the Universe, but in this paper there are no calculations confirming this fact.

There are other theories about 4th generation of fermions and mirror matter, for example:

- Sterile neutrino can be mirror particle. This neutrino haven't been observed by the present moment.
- All particles from the 4th generation of fermions are mirror.

etc. But in this paper these theories are not considered.

4. Literature.

[1] Lee T. D., Yang C. N. Phys. Rev. 104-254 (1956)

[2] I.Yu. Kobzarev, L.B. Okun, I.Ya. Pomeranchuk. On the possibility of observing mirror particles. Sov. J. Nucl. Phys. 3 (1966) 837

[3] Khlopov M.Yu., **Cosmoparticle Physics**

[4] <http://nuclphys.sinp.msu.ru/bm/bm06.htm>

[5] http://nuclphys.sinp.msu.ru/neutrino/newtrino_s/R&C.html

[6] B. Fields and S. Sarkar, "Big-Bang nucleosynthesis (2006, Particle Data Group mini-review),"

[7] **Cosmology**, Steven Weinberg, 2012

[8] J.A.Harvey, M.S. Turner, Phys. Rev. D 42, 3344 (1990)

[9] Zurab Berezhiani "Mirror world and its cosmological consequences", International Journal of Modern Physics, 2008

[10] Possible astronomical effects of mirror particles S. I. Blinnikov and M. Yu. Khlopov, Sov. J. Nucl. Phys. 36, 472 (1982)

[11] Wu C.S., Ambler E, Hayward R W, Hoppes D D, Hudson R P
«Experimental test of parity conservation in beta decay» , Physical Review,
vol.105, Issue 4, pp. 1413-1415, 1957

[12] Lee T. D., Yang C. N., Physical Review, vol. 104 (1): 254–258, 1955