

MINISTRY OF SCIENCE AND HIGHER EDUCATION OF THE RUSSIAN FEDERATION
FEDERAL STATE AUTONOMOUS EDUCATIONAL INSTITUTION HIGHER EDUCATION

"National Research Nuclear University "MEPhI"

Institute of Nuclear Physics and Technology

Small Thesis

"Quark-Gluon Blob in interaction of Ultra High Energy Cosmic Rays"

Student	<u>Abroo Uruj</u>
Group	<u>M21-192</u>
Lecturer	<u>M. Yu. Khlopov</u>

Moscow 2022

CONTENTS

1	Introduction.....	3
1.1	Unexplained events and corresponding requirement of new interaction model	4
2	Quark Gluon Hypothesis.....	5
3	QGM model explaining the observed anomalies	7
3.1	Calculations and comparison with experimental data	7
4	Evidences supporting QGM hypothesis.....	10
5	Conclusion	12
6	References.....	13


LIST OF FIGURES

Figure 1: Energy spectrum of cosmic rays.....	3
Figure 2: Non-central heavy-ion collisions with impact parameter $\sim b$. The global angular momentum of the produced matter is along $-\hat{y}$, opposite to the reaction plane.	6
Figure 3: Centrifugal barrier for particles with different masses: left – light mass, right – heavy mass.	6
Figure 4: Formation of measured cosmic ray energy spectrum in frame of nuclear-physical approach.	8
Figure 5: Changes of various nuclei spectra in the frame of the considered model.	9
Figure 6: Calculated and experimental data.....	9
Figure 7: Total orbital angular momentum of the overlapping system in Au+Au collisions at the RHIC energy as a function of the impact parameter b	10
Figure 8: Dependence of the magnitude of the excess of bundles of muons on the energy of primary cosmic rays for six existing models of interactions.....	11
Figure 9: Dependence of the average energy of muons in the bundles on the local muon density for the interval of zenith angles $\theta = 650 - 750$ [21].....	11

1 INTRODUCTION

High energy particles such as protons, photons and heavy nuclei, from extragalactic, galactic or solar origin are called cosmic rays. These primary cosmic rays, on entering the earth's atmosphere interact with nuclei in the atmosphere and produce what is called an air shower of secondary particles.

The global view of total energy spectrum of cosmic rays is as shown below:



The point on the spectrum at about $3 \cdot 10^{15}$ eV where the slope changes is called the knee and further, at about $3 \cdot 10^{18}$ eV, where the slope changes again is called the ankle. [1]

Observations of various phenomena relating to cosmic rays can be interpreted in either cosmological approach, where all changes are ascribed to changes in energy and/or composition of primary cosmic rays, or nuclear-physical approach,

i.e. based on the interaction model. Usually, the first approach is considered, where it is assumed that the energy of air showers is equal to that of the primary cosmic rays producing them. This perspective results in difficulties in explanation of many observations [2]. As is listed in the next section.

1.1 UNEXPLAINED EVENTS

At very high energies i.e. greater than 10^{15} eV, some observed events are inexplicable within the current theoretical purview [3][4]. The most puzzling characteristics of ultrahigh energy cosmic rays as observed in different experiments are listed below:

- In hadron experiments:
 - Families- Sets of separated cascades [5]
 - Halos around cascade families- diffused dark shadows around some cascade families
 - Alignment of cascades, despite the fact that probability of such random occurrence is negligible
 - Highly penetrating cascades
 - Centauros i.e. families of multiple hadron cascades without an accompanying electromagnetic cascade [6]
 - long-flying component that has large penetration before producing a cascade [7]
 - Anti-Centauros i.e. electromagnetic cascade without hadron cascade [8]
- In muon experiments:
 - Excess of very high energy (VHE~ 100 TeV) single (MSU) and multiple (LVD) muons[5]
 - Observation of VHE muons, even if the probability to detect such muons is very small. [9]
- In Extensive Air Shower (EAS) investigations: [10]
 - Young and old showers, large p_t , etc;
 - EAS spectrum change in atmosphere, usually interpreted as a change of primary energy spectrum;
 - $N_\mu(\text{Ne})$ and $X_{\text{max}}(\text{Ne})$ dependences change behaviour, explained as a change of the composition towards heavier elements.[11]

1.2 REQUIREMENT OF NEW INTERACTION MODEL

Therefore, the second approach i.e. nuclear-physical, can be considered instead. The main assumption of this alternate approach is that the energy of the EAS is not equal to the energy of the primary particles. Also, the new model of interaction that can explain all of the above listed observations would require to have the following characteristics:[3]

1. Threshold behaviour (since unusual events appear above several PeV only).
2. Large cross section (in order to change the slope of the EAS spectrum).
3. Large yield of leptons (to explain the excess of VHE muons and missing energy).
4. Large orbital (or rotational) momentum (to explain alignment).
5. Very fast development of EAS (for increasing N_{μ}/N_e ratio and decreasing X_{max} elongation rate).

2 QUARK GLUON HYPOTHESIS

All the above requirements are met if blobs of quark gluon matter with large orbital momentum are produced in non-central nucleus-nucleus collision that occur when high energy primary cosmic ray particles collide with the nuclei in the atmosphere at energies greater than several PeV [2][3]. It is important to note here, quark gluon matter implies partons in liquid state as opposed to plasma, which implies gas.

Quark gluon matter (QGM) production automatically meets the first requirement of threshold behaviour since production of QGM requires some high energy (temperature). Because of many quarks and gluons interacting in place of single pair of quark-quark interaction, geometrical cross section will be changed from $\sigma = \pi\lambda^2$ (for quark-quark interactions) to $\sigma = \pi(\lambda + R)^2$ or $\sigma \sim (R_1 + R_2)^2$ where R , R_1 , R_2 are sizes of interacting quark-gluon matter blobs. If R reaches even the size of nucleons, it will be sufficient to reach the necessary value of cross section.

Globally polarised QGM with large angular orbital momentum L is produced in non-central ion-ion collisions. L increases with increase in collision energy in center of mass system $L \sim \sqrt{s}$ [12].




Figure 2: Non-central heavy-ion collisions with impact parameter $\sim b$. The global angular momentum of the produced matter is along $-\hat{y}$, opposite to the reaction plane.

The centrifugal barrier of the QGM increases with L and thus with the collision energy \sqrt{s} . The blob here can be considered to be in resonance state with increased lifetime, with a large centrifugal barrier. This centrifugal force has a value in centre of mass system of $V(L) = L^2/mr^2$. Thus, the centrifugal barrier will be lower for heavier quarks and the probability of its decay will be larger. This results in the suppression of light quark production, and thus provides enough time for appearance of heavy t quarks (or other heavy particles) in the very hot QGM. [13]



Figure 3: Centrifugal barrier for particles with different masses: left – light mass, right – heavy mass.

This top quark (or any other heavy particle with mass $\gtrsim 100 \text{ GeV}/c^2$, taking into account fly out energy) will decays into W or Z boson.

$$\text{eg: } t(\bar{t}) \rightarrow W^+(W^-) + b(\bar{b}) \quad [14]$$

Here W - boson and

b - quark

3 QGM MODEL EXPLAINING THE OBSERVED ANOMALIES

The bosons thus produced will decay into hadrons (probability $\approx 70\%$, on average, and $\approx 30\%$ pions) which explains appearance of excess of muon bundle, due to multiplicity of the secondary particles produced [3]. Correspondingly EAS development varies which can explain young and old showers, large values of transverse momentum, etc.

Alternatively, decay of the W bosons into lepton pairs ($e\nu_e, \mu\nu_\mu, \tau\nu_\tau$ with probability $\approx 10\%$ per each pair) can account for the missing energy as these neutrinos are not detected and the muon energy is not measured either. The energy of these muons and neutrinos will be very large (more than 100 TeV) and thus the measured energy of EAS can sufficiently differ from the primary particle energy, causing a knee in measured spectrum.

Penetrating cascades can be explained too. VHE muons can produce similar events. The flux of VHE muons from known sources with necessary energies (~ 1 PeV) is very small. However, muons from decays of heavy particles (through W, Z-bosons) increase the flux at these energies by at least two orders

Centauros and other such unusual events' probability is increased, as in weak interactions, there is non-conservation of isospin and only charged particle production will be possible.

3.1 CALCULATIONS AND COMPARISON WITH EXPERIMENTAL DATA

Collision energy changes because of simultaneous interaction of many quarks:

$$\sqrt{s} = \sqrt{2m_N E_1} \rightarrow \sqrt{2m_C E_1}$$

where m_N = nucleon mass and

m_C is compound mass of many interacting quarks.

For calculations, one can consider $m_C = nm_N$, ($n = 1 \div A$).

Production of $t\bar{t}$ -quark pair must decrease \sqrt{s} at least by the value of $2m_t$, and in a general case by some value $\varepsilon_t > 2m_t \approx 350 GeV$ which will depend on primary particle energy and its mass. The residual part of the energy in the center

of mass system $(\sqrt{s} - \varepsilon_t)$ will be converted into the energy of usual processes of EAS development. Some part of the energy taken away by top-quarks will be re-injected into EAS development, but in the first approximation, to simplify consideration, it is possible to neglect this value. So, results of standard measurements and standard procedure of evaluation of EAS energy will give value

$$E_2 = \frac{(\sqrt{2m_c E_1} - \varepsilon_t)^2}{m_c}$$

which is less than the energy of the primary particle E_1 , and we will obtain the steepening of the observed spectrum

Transition from E_1 to E_2 gives a bump near the threshold as shown in figure below [2]




Figure 4: Formation of measured cosmic ray energy spectrum in frame of nuclear-physical approach.

As high density is required for the production of QGM, the threshold energy for production of new state of matter will be lesser for heavier nuclei compared to that for lighter nuclei. Thus, it is assumed that at first QGM is produced in iron nuclei (or any other heavier nuclei) interactions. Following this increasingly lighter nuclei produce the QGM. This is shown in the figure below:




Figure 5: Changes of various nuclei spectra in the frame of the considered model.

For calculations primary spectra values for various nuclei were taken from [15]. The results of calculations assuming 15% straggling of measured energies and comparison with experimental data on all particle spectrum are given in Figure.




Figure 6: Calculated and experimental data.

4 EVIDENCES SUPPORTING QGM HYPOTHESIS

At the Relativistic Heavy Ion collider (RHIC), during Au-Au collisions at $\sqrt{s} = 200$ GeV, the total angular orbital momentum L of the overlapping system as a function of impact parameter b , obtained is shown in the figure. [13], [16] Also shown in the plot, is the curve in dashed line for a hard sphere (which will not gain a large angular momentum upon non central impact). Comparing the two curves it becomes clear that the postulate about increased Woods - Saxon potential with increase in momentum for heavier particles is true.




Figure 7: Total orbital angular momentum of the overlapping system in Au+Au collisions at the RHIC energy as a function of the impact parameter b .

Upon simulation of EAS cascade while including the production of top-pair in its development, excess of muons with energy ~ 100 TeV appears, if the production of blobs of QGM starts at energies of several PeV[14]. Thus, potentially resolving muon puzzles.

Practically, this hypothesis can be directly proven by either accurately measuring the energy spectrum of atmospheric muons above energies of 100TeV or by showing that excess of VHE muons appear in interactions of primary particles with energy above the knee and are absent below the knee. [3].

The dependence of the excess of muon bundles on the energy of primary cosmic rays, for six existing interaction models mentioned in the graphs, that were registered by all the different installations that registered multiple muons generated by primary cosmic rays of ultrahigh energies is as shown below[[17]. The largest contribution in the energy range from 10^{14} to 10^{20} eV was made by

three installations: NEVOD, Pierre Auger, and IceCube. The experiment at MEPHI covers the largest energy range, in the lower energy region it is proved by the IceCube data and by Pierre Auger in the ultrahigh energy region.




Figure 8: Dependence of the magnitude of the excess of bundles of muons on the energy of primary cosmic rays for six existing models of interactions.

According to the data of measurements of bundles of muons at the NEVOD-DECOR complex [18][19][20] in the range of zenith angles of $65^\circ - 75^\circ$, corresponding to the energies of primary particles above 10^{17} eV, an increase in the average energy of muons in bundles was found in comparison with the results of calculations using modern post-LHC models of hadronic interactions, which may indicate a change in the interaction model [21].




Figure 9: Dependence of the average energy of muons in the bundles on the local muon density for the interval of zenith angles $\theta = 65^\circ - 75^\circ$ [21].

The inclusion of a new coordinate-track detector TREK in the NEVOD experimental complex will allow to expand the range of investigated energies to the interval from 10^{14} to 10^{19} eV [25], which will cover the entire region in which an excess of cosmic ray muons is observed. To improve the accuracy of muon bundle energy deposit measurement, the extension and optimization of Cherenkov water detector structure will be done [26]. An increase in the accuracy in determining the multiplicity of bundles and their energy deposit will make it possible to understand the nature of the muon puzzle and give an answer about the inclusion of the QGM hypothesis in the interaction of primary cosmic rays of ultrahigh energies.

5 CONCLUSION

Anomalies observed in ultrahigh energy cosmic ray spectrum are discussed.

Hypothesis of production of Quark Gluon Blobs when ultrahigh energy cosmic ray particles collide with particles in the atmosphere and the consequences following it are detailed.

Experimental results that support the postulates and that experimental setup that can prove this hypothesis is discussed.

6 REFERENCES

- [1] T. K. Gaisser, R. Engel, and E. Resconi, *Cosmic {Rays} and {Particle} {Physics}*, 2nd ed. Cambridge: Cambridge University Press, 2016. doi: 10.1017/CBO9781139192194.
- [2] A. A. Petrukhin, 'Cosmic rays above the knee: experimental results and their interpretation', p. 4.
- [3] A. A. Petrukhin, 'Manifestations of quark-gluon plasma in cosmic ray experiments', *Nuclear Physics B - Proceedings Supplements*, vol. 175–176, pp. 125–128, Jan. 2008, doi: 10.1016/j.nuclphysbps.2007.10.020.
- [4] V. M. Maximenko, V. S. Puchkov, S. E. Pyatovsky, S. A. Slavatinsky, and R. A. Mukhamedshin, '{SOME} {INTERESTING} {PHENOMENA} {OBSERVED} {IN} {COSMIC}-{RAY} {EXPERIMENTS} {BY} {MEANS} {OF} {X}-{RAY} {EMULSION} {TECHNIQUE} {AT} {SUPER} {ACCELERATOR} {ENERGIES}', p. 16.
- [5] A. A. Petrukhin, 'muon excess.pdf'. *Conference Proceedings Vol. 85, 'Frontier Objects in Astrophysics and Particle Physics'* P.527, p. 12, 2003.
- [6] E. Gladysz-Dziadus, 'Are {Centaurus} exotic signals of the {QGP}?', arXiv:hep-ph/0111163, Jan. 2012.
- [7] A. O. Herrera et al., '{EDITOR} {DA} {DIVERSIDADE} {ESTADUAL} {DE} {CAMPINAS} {Reitor}: {J} ». -• {Aristodemo} {Pinotti}', p. 260.
- [8] V. I. Yakovleva, 'Anti-{Centaur0} type events at energies 5-500 {TeV} detected by the {Tien}-{Shan} {EAS} complex', p. 1.
- [9] E. Iarocci and S. Ragazzi, 'Cosmic-Muon Results from the NUS] ~ X Experiment .', 1986.
- [10] S. B. Shaulov et al., 'Investigation of EAS cores', *EPJ Web of Conferences*, vol. 145, no. November, p. 17001, 2017, doi: 10.1051/epjconf/201714517001.
- [11] A. Aab et al., 'Depth of maximum of air-shower profiles at the Pierre Auger Observatory. I. Measurements at energies above $>1 ⁰ </math> $>17.8 </math> $<mi>eV</mi>$ ', *Physical Review D*, vol. 90, no. 12, p. 122005, 2014, doi: 10.1103/PhysRevD.90.122005.$$
- [12] Z. T. Liang and X. N. Wang, 'Globally polarized quark-gluon plasma in noncentral A + A collisions', *Physical Review Letters*, vol. 94, no. 10, pp. 1–5, 2005, doi: 10.1103/PhysRevLett.94.102301.
- [13] J. H. Gao, S. W. Chen, W. T. Deng, Z. T. Liang, Q. Wang, and X. N. Wang, 'Global quark polarization in noncentral A+A collisions', *Physical Review C - Nuclear Physics*, vol. 77, no. 4, pp. 1–13, 2008, doi: 10.1103/PhysRevC.77.044902.
- [14] A. A. Petrukhin, 'Muon puzzle in cosmic ray experiments and its possible solution', *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, vol. 742, pp. 228–231, Apr. 2014, doi: 10.1016/j.nima.2013.12.011.
- [15] P. D. Group, 'University of Zurich', *Journal of Clinical Epidemiology*, vol. 58, no. 5, p. I, 2005, doi: 10.1016/s0895-4356(05)00103-4.
- [16] Z.-T. Liang and X.-N. Wang, 'Globally Polarized Quark-gluon Plasma in Non-central A+A Collisions', *Phys. Rev. Lett.*, vol. 94, no. 10, p. 102301, Mar. 2005, doi: 10.1103/PhysRevLett.94.102301.
- [17] E. A. Zadeba et al., 'The registration system of the coordinate-tracking setup on the drift chambers', *J. Phys.: Conf. Ser.*, vol. 675, no. 3, p. 032039, Feb. 2016, doi: 10.1088/1742-6596/675/3/032039.
- [18] Dembinski H P 2019 The Muon Puzzle in High-Energy Air Showers. *Phys. At. Nucl.* 82 644.

- [19] Bogdanov A G et al. 2018 Investigation of very high energy cosmic rays by means of inclined muon bundles. *Astropart. Phys.* 98 13.
- [20] Bogdanov A G et al. 2020 Intensity of muon bundles according to the NEVOD-DECOR cosmic ray experiment *Journal of Physics: Conference Series* 1690(1) 012007.
- [21] Yurina E A et al. 2019 Investigation of the energy loss of muon bundles in the Cherenkov water calorimeter *Physics of Atomic Nuclei* 82 680–684.
- [22] Zadeba E A et al. 2014 The coordinate-tracking detector based on the drift chambers for ultrahigh-energy cosmic ray investigations *Journal of Instrumentation* 9 C08018.
- [23] Zadeba E A et al. 2017 The detector on the basis of drift chambers for inclined muon bundle investigations *Journal of Instrumentation* 12 75.
- [24] Zadeba E A et al. 2019 Investigation of muon bundles generated by UHECR by means of the new coordinate-tracking detector *Journal of Physics: Conference Series* 1390(1) 012132.
- [25] Khokhlov S S et al. 2021 Studying the Characteristics of Optical Modules in the Volume of the NEVOD Cherenkov Water Detector *Bull. RAS: Phys* 85.4 452-454.