

Particle - Mirror Particle Oscillations: *CP-violation and Baryon Asymmetry*

Zurab Berezhiani

Università di L'Aquila and LNGS, Italy

'Now, if you'll only attend, Kitty, and not talk so much, I'll tell you all my ideas about Looking-glass House. There's the room you can see through the glass – that's just the same as our drawing-room, only the things go the other way... the books are something like our books, only the words go the wrong way: I know that, because I've held up one of our books to the glass, and then they hold up one in the other room. I can see all of it – all but the bit just behind the fireplace. I do so wish I could see that bit! I want so to know whether they've a fire in the winter: you never can tell, you know, unless our fire smokes, and then smoke comes up in that room too – but that may be only pretence, just to make it look as if they had a fire...

'How would you like to leave in the Looking-glass House, Kitty? I wonder if they'd give you milk in there? But perhaps Looking-glass milk isn't good to drink? Now we come to the passage: it's very like our passage as far as you can see, only you know it may be quite on beyond. Oh, how nice it would be if we could get through into Looking-glass House! Let's pretend there's a way of getting through into it, somehow ... Why, it's turning into a sort of mist now, I declare! It'll be easy enough to get through ...'

–Alice said this, and in another moment she was through the glass... she was quite pleased to find that there was a real fire in the fireplace... 'So I shall be as warm here as I was in my room,' thought Alice: 'warmer, in fact, there'll be no one here to scold me away from the fire.'

● Carroll's Alice...

- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

"Looking-Glass Universe" – Parallel "Mirror" World

Broken P can be restored by **mirror fermions**

Lee & Yang '56

Mirror sector **hidden copy** of our sector

Kobzarev, Okun, Pomeranchuk '66

Alice strings

A.S. Schwarz' ??

Mirror dark matter (invisible stars)

Blinnikov, Khlopov '83

$$SU(3) \times SU(2) \times U(1) \quad \times \quad SU(3)' \times SU(2)' \times U(1)'$$

Foot, Lew, Volkas '91

Two identical gauge factors, $G \times G'$, with the identical field contents and Lagrangians: $\mathcal{L}_{\text{tot}} = \mathcal{L} + \mathcal{L}' + \mathcal{L}_{\text{mix}} \quad - \quad SU(5) \times SU(5)', \text{ etc.}$

- Can naturally emerge in string theory: O & M matter fields localized on two parallel branes with gravity propagating in bulk: e.g. $E_8 \times E_8'$

- Exact parity $G \leftrightarrow G'$: Mirror matter is dark (for us), but its particle physics we know exactly – **no new parameters!**

- Spont. broken parity $G \leftrightarrow G'$: $M'_W \neq M_W \quad (M'_W \geq \text{few TeV})$

Particle spectrum rescaled by $\zeta = M'_W/M_W$

ZB & Mohapatra '95

Shadow DM, sterile neutrinos, Machos

ZB, Dolgov, Mohapatra '96

Strong CP and new axion (axidragon)

ZB, Gianfagna, Giannotti '00

SUSY little Higgs – accidental global $U(4)$

ZB '04

- Carrol's Alice...
- **Mirror World**
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Mirror Sector, Mirror Particles & Mirror Parity

$SU(3) \times SU(2) \times U(1)$
gauge (g, W, Z, γ)
& Higgs (ϕ) fields

$SU(3)' \times SU(2)' \times U(1)'$
gauge (g', W', Z', γ')
& Higgs (ϕ') fields

quarks ($B=1/3$)	leptons ($L=1$)		quarks ($B'=1/3$)	leptons ($L'=1$)
$q_L = (u, d)_L^t$	$l_L = (\nu, e)_L^t$		$q'_L = (u', d')_L^t$	$l'_L = (\nu', e')_L^t$
$u_R \quad d_R$	e_R		$u'_R \quad d'_R$	e'_R
$\widetilde{\text{quarks}} (B=-1/3)$	$\widetilde{\text{leptons}} (L=-1)$		$\widetilde{\text{quarks}} (B'=-1/3)$	$\widetilde{\text{leptons}} (L'=-1)$
$\tilde{q}_R = (\tilde{u}, \tilde{d})_R^t$	$\tilde{l}_R = (\tilde{\nu}, \tilde{e})_R^t$		$\tilde{q}'_R = (\tilde{u}', \tilde{d}')_R^t$	$\tilde{l}'_R = (\tilde{\nu}', \tilde{e}')_R^t$
$\tilde{u}_L \quad \tilde{d}_L$	\tilde{e}_L		$\tilde{u}'_L \quad \tilde{d}'_L$	\tilde{e}'_L

$$- \mathcal{L}_{\text{Yuk}} = f_L Y \tilde{f}_L \phi + \tilde{f}_R Y^* f_R \tilde{\phi} \quad | \quad \mathcal{L}'_{\text{Yuk}} = f'_L Y' \tilde{f}'_L \phi' + \tilde{f}'_R Y'^* f'_R \tilde{\phi}'$$

- D-parity: $L \leftrightarrow L', R \leftrightarrow R', \phi \leftrightarrow \phi' : Y' = Y$ • *identical xero copy*
- M-parity: $L \leftrightarrow R', R \leftrightarrow L', \phi \leftrightarrow \tilde{\phi}' : Y' = Y^\dagger$ • *mirror (chiral) copy*

- Carroll's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Possible interactions between O & M particles (besides gravity)

(but also model of mirror gravity can be constructed !)

Can be at tree level, or induced by exchange of extra gauge singlet particles or common gauge fields acting with both O & M particles ...

another interesting story !

Z. Berezhiani, *Phys. Lett. B* 417, 287 (1998)

*these interactions can induce particle mixing phenomena between O & M sectors:
any neutral particle (elementary or composite) can mix its mirror twin
exactly degenerate in mass*

■ photon - mirror photon kinetic mixing $\epsilon F^{\mu\nu} F'_{\mu\nu}$ Holdom '86

mirror particles become "millicharged" $Q' \sim \epsilon Q$ relative to our photon

→ positronium - mirror positronium mixing ($e^+ e^- \rightarrow e'^+ e'^-$) Glashow '86

... but BBN : $\epsilon < 10^{-8}$, CMB+LSS : $\epsilon < 10^{-9}$

■ meson - mirror meson mixing: $\pi^0 - \pi^{0'}$, $K^0 - K^{0'}$, $\rho^0 - \rho^{0'}$, etc.

$\frac{1}{M^2} (\bar{u}\gamma^5 u - \bar{d}\gamma^5 d)(\bar{u}'\gamma^5 u' - \bar{d}'\gamma^5 d')$, $\frac{1}{M^2} (\bar{d}\gamma^5 s)(\bar{d}'\gamma^5 s')$ ($\Delta S = 1$)

... analogous to $\frac{1}{M^2} (\bar{d}\gamma^5 s)(\bar{d}\gamma^5 s) \rightarrow K^0 - \bar{K}^0$ mixing ($\Delta S = 2$)

Phenom. limits: $M > 10 \text{ TeV}$ ($\pi^0 - \pi^{0'}$), $M > 100 \text{ TeV}$ ($K^0 - K^{0'}$)

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Lepton & baryon number violating interactions

- neutrino - mirror neutrino mixing ($\nu - \nu'$) – effective operators :
Z. Berezhiani, R.N. Mohapatra, Phys. Rev. D 52, 6607 (1995)

$$\frac{1}{M} (l\phi)(l'\phi') \quad (\Delta L = 1, \Delta L' = 1)$$

analogous to $\frac{1}{M} (l\phi)^2 \quad (\Delta L = 2), \quad \frac{1}{M} (l'\phi')^2 \quad (\Delta L' = 2)$

– operators that generate neutrino Majorana masses via seesaw mechanism
 constraints from active-sterile neutrino mixing

- neutron - mirror neutron mixing ($n - n'$) – effective operators :

$$\frac{1}{M^5} (udd)(u'd'd'), \quad (\Delta B = 1, \Delta B' = 1)$$

analogous operators $\frac{1}{M^5} (udd)^2 \quad (\Delta B = 2), \quad \frac{1}{M^5} (u'd'd')^2 \quad (\Delta B' = 2)$

generate neutron - antineutron mixing

- hydrogen - mirror hydrogen mixing – effective operators :

$$\frac{1}{M^8} (udde)(u'd'd'e'), \quad (\Delta B = 1, \Delta L = 1; \Delta B' = 1, \Delta L' = 1)$$

c.f. operators $\frac{1}{M^8} (udde)^2 \longrightarrow$ *hydrogen - antihydrogen atom mixing*

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

BBN demands : was Alice's guess correct?

Mirror particle physics \equiv ordinary particle physics

but mirror cosmology \neq ordinary cosmology

■ at the BBN epoch, $T \sim 1 \text{ MeV}$, $g_* = g_*^{SM} = 10.75$
as contributed by the γ , e^\pm and 3ν species : $N_\nu = 3$

■ if $T' = T$, mirror world would give the same contribution:

$$g_*^{\text{eff}} = 2 \times g_*^{SM} = 21.5 \text{ – equivalent to } \Delta N_\nu = 6.14 \text{ !!!}$$

■ If $T' < T$, then $g_*^{\text{eff}} \approx g_*^{SM} (1 + x^4)$, $x = T'/T \longrightarrow \Delta N_\nu = 6.14 \cdot x^4$
E.g. $\Delta N_\nu < 0.4$ requires $x < 0.5$; for $x = 0.2$ $\Delta N_\nu \simeq 0.01$

■ Paradigm – different initial conditions & weak contact :

– after inflation O and M worlds are (re)heated non-symmetrically, $T' < T$

– processes between O - M particles are slow enough & stay **Out-of-Equilibrium**

– both sectors evolve **adiabatically**, without significant entropy production

So $x = T'/T$ is nearly independent of time ($T'_{\text{CMB}}/T_{\text{CMB}}$ today)

BBN: $\Delta N_\nu/6.14 = x^4 \ll 1 \longrightarrow$ BBN': $\Delta N'_\nu/6.14 = x^{-4} \gg 1$

^1H 75%, ^4He 25% vs. $^1\text{H}'$ 25%, $^4\text{He}'$ 75%

Z. Berezhiani, D. Comelli, F. Villante, Phys. Lett. B 503, 362 (2001)

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Cosmic Coincidence & Fine Tuning Problems

Today's Universe is flat ($\Omega_{\text{tot}} \approx 1$) and multi-component:

- $\Omega_B \simeq 0.04$ observable matter – Baryons !
- $\Omega_D \simeq 0.20$ dark matter: – WIMPS? Axions?
- $\Omega_\Lambda \simeq 0.75$ dark energy: – Λ -term? 5th-essence?

A. coincidence of matter $\Omega_M = \Omega_D + \Omega_B$ and dark energy Ω_Λ : $\Omega_M / \Omega_\Lambda \simeq 0.3$

· $\rho_\Lambda \sim \text{Const.}$, $\rho_M \sim a^{-3}$; why $\rho_M / \rho_\Lambda \sim 1$ – just Today?

Anthropic answer: if not Today, then it could be Yesterday or Tomorrow ...

B. Fine Tuning between visible Ω_B and dark Ω_D matter: $\Omega_B / \Omega_D \simeq 0.2$

· $\rho_B \sim a^{-3}$, $\rho_D \sim a^{-3}$; why $\rho_B / \rho_D \sim 1$ – Yesterday Today & Tomorrow?

Difficult question ... popular models for the primordial Baryogenesis (GUT-B, Lepto-B, Spont. B, Affleck-Dine B, EW B, ...) have no feeling for the popular DM candidates (Wimp, Wimpzilla, axion, axino, gravitino ...)

– *How Baryon Asymmetry could know about Dark Matter?* – again anthropic (landscaped) Fine Tunings in Particle Physics and Cosmology? Just for our good?

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Visible vs. Dark matter

- Visible matter: $\rho_B = n_B M_B$, $M_B \simeq 1 \text{ GeV}$ – nucleons, $\eta = n_B/n_\gamma \sim 10^{-9}$

Sakharov's conditions: B ($B - L$) & CP violation, Out-of-Equilibrium

– in Baryogenesis models η depends on several factors, like CP-violating constants, particle degrees of freedom, mass scales, particle interaction strength and goodness of out-of-equilibrium.... and in some models (e.g. Affleck-Dine) on the initial conditions as well...

- Dark matter: $\rho_D = n_X M_X$, but $M_X = ?$, $n_X = ?$

– too wide spectrum of possibilities ...


Axion: $M_X \sim 10^{-5} \text{ eV}$; **Wimp:** $M_X \sim 1 \text{ TeV}$; **Wimpzilla:** $M_X \sim 10^{14} \text{ GeV}$...

– in relative models n_X depends on various factors, like equilibrium status and particle degrees of freedom, particle masses and interaction strength (production and annihilation cross sections).... and in some models (e.g. Axion or Wimpzilla) on the initial conditions as well ...

How then the mechanisms of Baryogenesis and Dark Matter synthesis, having different particle physics and corresponding to different epochs, could know about each-other? – How $\rho_B = n_B M_B$ could match $\rho_X = n_X M_X$ so intimately?

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment


B vs. D – Fine Tuning demonstration



Evolution of the Baryon number (\cdots) in e.g. Leptogenesis scenario confronted to the evolution of the Dark Matter density ($-$) in the scenarios of WIMP (left pannel) and Axion (right pannel)

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- **B vs. D – Fine Tuning demonstration**
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Unified origin of B and D? Both fractions at one shoot?



$$\frac{\rho_X}{\rho_B} = \frac{M_X n_X}{M_B n_B} \sim 1 \quad \text{can be natural}$$

- if DM properties are similar to baryon ones: namely $M_X \sim M_B$
- and both fractions are generated by same mechanism so that $n_X \sim n_B$

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- **Unification**

- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

O & M neutrino mixing

Mixed $D=5$ effective operators

Z.B. & Mohapatra '95

$$\frac{A}{M} ll\phi\phi_{(\Delta L=2)} + \frac{A'}{M} l'l'\phi'\phi'_{(\Delta L'=2)} + \frac{D}{M} ll'\phi\phi'_{(\Delta L=1, \Delta L'=1)}$$

Substituting VEVs $\langle\phi\rangle = v$ and $\langle\phi'\rangle = v'$, we get $\nu - \nu'$ mixing

$$\begin{pmatrix} \hat{m}_\nu & \hat{m}_{\nu\nu'} \\ \hat{m}_{\nu\nu'}^t & \hat{m}_{\nu'} \end{pmatrix} = \frac{1}{M} \begin{pmatrix} Av^2 & Dvv' \\ D^t vv' & A'v'^2 \end{pmatrix} - \text{active-sterile } \nu \text{ system}$$

[M-parity: $A' = A^*$, $D = D^\dagger$; D-parity: $A' = A$, $D = D^t$]

• $v' = v$: $m_{\nu\nu'} = m_\nu$ and maximal mixing $\theta_{\nu\nu'} = 45^\circ$; Foot & Volkas '95

• $v' > v$: $m_{\nu\nu'} \sim (v'/v)^2 m_\nu$ and small mixing $\theta_{\nu\nu'} \sim v/v'$;

e.g. $v'/v \sim 10^2$: $\sim \text{keV sterile neutrinos as WDM}$ Z.B., Dolgov, Mohapatra '96

• $A, A' = 0$ ($L-L'$ conserved) light – Dirac neutrinos Z.B. & Bento '05
with L components in ordinary sector and R components in mirror sector

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Mixed Seesaw and Leptogenesis between O & M sectors

- Heavy gauge singlet fermions N_a , $a = 1, 2, 3, \dots$ with large Majorana mass terms $M_{ab} = g_{ab}M$, can equally talk with both O and M leptons

$$\mathcal{L}_{\text{Yuk}} = y_{ia}\phi l_i N_a + y'_{ia}\phi' l'_i N_a + \frac{1}{2}Mg_{ab}N_a N_b + \text{h.c.};$$

(M-parity: $y' = y^\dagger$; D-parity: $y' = y$)

- D=5 effective operators $\frac{A}{M}ll\phi\phi + \frac{A'}{M}l'l'\phi'\phi' + \frac{D}{M}ll'\phi\phi'$ emerge after integrating out heavy states N , where

$$A = yg^{-1}y^t, \quad A' = y'g^{-1}y'^t, \quad D = yg^{-1}y'^t$$

- They generate also processes like $l\phi \rightarrow \tilde{l}'\tilde{\phi}'(l'\phi')$ ($\Delta L = 1$) and $l\phi \rightarrow \tilde{l}\tilde{\phi}$ ($\Delta L = 2$) satisfying Sakharov's 3 conditions for baryogenesis

A. violate B-L – *by definition*

B. violate CP – *complex Yukawa constants y_{ia}*

C. out-of-equilibrium – *already implied by the BBN*

and thus generate $B-L \neq 0$ ($\rightarrow B \neq 0$ by sphalerons) for ordinary matter

- The same reactions generate $B'-L' \neq 0$ ($\rightarrow B' \neq 0$) in Mirror sector.

Both matter fractions: observable and dark, can be generated at one shoot !!


- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing

● See-Saw

- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

CP violation in $\Delta L=1$ and $\Delta L=2$ processes

L. Bento, Z. Berezhiani, PRL 87, 231304 (2001)



$$\varepsilon_{CP} = \text{Im Tr}[(y^\dagger y)^* g^{-1} (y'^\dagger y') g^{-2} (y^\dagger y) g^{-1}]$$

$$\varepsilon'_{CP} = \text{Im Tr}[(y'^\dagger y')^* g^{-1} (y^\dagger y) g^{-2} (y'^\dagger y') g^{-1}]$$

$$\varepsilon_{CP} \rightarrow \varepsilon'_{CP}$$

when $y \rightarrow y'$


- **D-parity:** $y' = y$, $\varepsilon_{CP} = 0$, but **M-parity:** $y' = y^\dagger$ $\varepsilon_{CP} \neq 0$

- Carroll's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- **Leptogenesis: diagrams**
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Leptogenesis: $M'_B = M_B \dots$ but $\Omega'_B \geq \Omega_B$

$$B = D(k) \cdot Y^{(0)}, \quad B' = D(kx^3) \cdot Y^{(0)}; \quad Y^{(0)} \approx \frac{\varepsilon_{CP} M_{Pl} T_R^3}{g_*^{3/2} M^4} \cdot 10^{-3}$$

$$k = [\Gamma_{\text{eff}}/H]_{T=T_R}, \quad x = T'/T \approx 1.2 (k/g_*)^{1/4} \quad (T_R = T_{\text{Reheating}})$$



Z.B. '03

BBN: $x < 0.5 \rightarrow k \leq 4$; LSS: $x < 0.2 \rightarrow k \leq 1.5$

Thus Ordinary/Mirror matter ratio can vary within $\frac{\Omega_B}{\Omega'_B} = D(k) \simeq 0.2 - 1$

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- **Leptogenesis: formulas**
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Mirror Baryons as Dark Matter

As far as Mirror Baryons are dark (in terms of ordinary photons), they could constitute Dark Matter of the Universe [Z.B., Comelli & Villante '01]

- Once $x < 1$, mirror photons decouple earlier than our photons: $z'_{\text{dec}} \simeq \frac{1}{x} z_{\text{dec}}$

However, if the DM is entirely due to mirror baryons, then the large scale structure (LSS) formation requires that mirror photons must decouple before Matter-Radiation Equality epoch: $x < x_{\text{eq}} = 0.05(\Omega_M h^2)^{-1} \simeq 0.3$

- then mirror Jeans scale λ'_J becomes smaller than the Hubble horizon before Matter-Radiation Equality

- mirror Silk scale is smaller than the one for the normal baryons:

$$\lambda'_S \sim 5x_{\text{eq}}^{5/4} (x/x_{\text{eq}})^{3/2} (\Omega_M h^2)^{-3/4} \text{ Mpc}$$

Hence the structures formation at 1 Mpc scales (galaxies) implies $x < 0.2$



N.B. Since mirror baryons constitute dissipative dark matter, the formation of the extended halos can be problematic, but perhaps possible if the star formation in the mirror sector is rather fast due to different temperature and chemical content (in fact, fast freezout of BBN in mirror sector is much faster, and it is dominated by Helium).

MACHOs as mirror stars – microlensing: $M_{\text{av}} = 0.5 M_{\odot}$

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Leptogenesis or Baryogenesis?

$D=5$ operator $\frac{1}{M} ll' \phi \phi'$ ($\Delta L = 1$) induced by heavy singlet N "seesaw" exchange (l, ϕ and l', ϕ' ordinary and mirror lepton and Higgs doublets) – can generate $B-L$ (and $B' - L'$) asymmetry via processes $l\phi \rightarrow l'\phi'$ Z.B. and Bento '01



Z.B. and Bento '05

$D=9$ operator $\frac{1}{M^5} (udd)(u'd'd')$ ($\Delta B = 1$) induced by heavy singlet N "seesaw" (u, d and u', d' ordinary and mirror R -quarks, S, S' color triplet scalars (squarks?)) – can generate $B-L$ (and $B' - L'$) asymmetry via processes $dS \rightarrow d'S'$

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Neutron - Mirror neutron mixing

Operators like $\frac{1}{\mathcal{M}^5} (udd)(u'd'd')$ and $\frac{1}{\mathcal{M}^5} (qqd)(q'q'd')$ induce the neutron - mirror neutron mass mixing $\delta m (\bar{n}n' + \bar{n}'n)$, with $\delta m \sim \left(\frac{10 \text{ TeV}}{\mathcal{M}}\right)^5 \cdot 10^{-15} \text{ eV}$

• $n - n'$ oscillation in vacuum:

maximal mixing $\theta = 45^\circ$ and oscillation time $\tau_{\text{osc}} = \delta m^{-1} \sim \left(\frac{\mathcal{M}}{10 \text{ TeV}}\right)^5 \text{ s}$

... similar to neutron - antineutron oscillation

Kuzmin '70, Glashow '79

Marshak & Mohapatra '80

but experimental limits on $n - \bar{n}$ are strong: $\tau_{n\bar{n}} > 10 \text{ yr}$, while $n - n'$ is still allowed to be rather fast, faster than neutron decay: $\tau_{nn'} < 10 \text{ min}$

Can be interesting if $\mathcal{M} \sim (M_S^4 M_N)^{1/5} \sim 10 \text{ TeV}$ In the "seesaw" model – E.g. if $M_S, M_N \sim 10 \text{ TeV}$, or $M_N \sim 10^{12} \text{ TeV}$ and $M_S \sim 100 \text{ GeV}$

(see diagram of the previous page)

!!! N.B. Nuclear Stability

• $n - \tilde{n}$ destabilizes nuclei: $(A, Z) \rightarrow (A - 1, Z, \tilde{n}) \rightarrow (A - 2, Z) + \pi$'s

$\tau_{n\tilde{n}} > 10 \text{ yr}$ or so ...

• $n - n'$ does not: $(A, Z) \rightarrow (A - 1, Z) + n'$ **not allowed by phase space !**
gives no restriction for $\tau_{nn'}$!

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Neutron - Mirror neutron oscillation in external fields

Effective (non-relativistic) Hamiltonian for $n - n'$ oscillation

Z.B. & Bento '05

$$H = \begin{pmatrix} m - i\Gamma/2 + V + \mu(\vec{B} \cdot \vec{\sigma}) & \delta m \\ \delta m & m' - i\Gamma'/2 + V' + \mu'(\vec{B}' \cdot \vec{\sigma}) \end{pmatrix}$$

- Exact mirror parity: $m' = m, \Gamma' = \Gamma, \mu' = \mu = -1.91\mu_N$
- Grav. potentials are the same: $V' = V,$
- but magnetic fields $\vec{B}' \neq \vec{B}$:

$$|\mu B| \simeq 6 \cdot 10^{-12} \text{ eV/G} \quad (\text{Earth magnetic field } B \simeq 0.5 \text{ G}) -$$

Take $\mathbf{B} = (0, 0, B)$ across z -axis, $(\boldsymbol{\sigma} \mathbf{B}) = B\sigma_z = \text{diag}(B, -B)$ and $B' = 0$

$$H = \begin{pmatrix} \pm 2\omega_B & \delta m \\ \delta m & 0 \end{pmatrix} \quad \text{diagonal in the basis } (\psi_+, \psi_-, \psi'_+, \psi'_-)$$

– Energy gap $2\omega_B = |\mu B| \simeq B[\text{G}] \times 6 \cdot 10^{-12} \text{ eV}$

Oscillation probability $P_{nn'}(t) = \sin^2 2\theta_B \sin^2(t/\tau_B) \cdot e^{-t/\tau_{\text{dec}}}$

$$\sin 2\theta_B = \frac{\delta m}{\sqrt{\delta m^2 + \omega_B^2}}, \quad \tau_B = \frac{1}{\sqrt{\delta m^2 + \omega_B^2}} = \tau \sin 2\theta_B, \quad \tau = \delta m^{-1}$$

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

$n - n'$ transition probabilities ($B' = 0, t \ll \tau_{\text{dec}}$)

In vacuum ($\omega_B = 0$): $P_0(t) = \sin^2(\delta m t)$

($\tau_0 = \tau = \delta m^{-1}, \sin^2 2\theta_0 = 1 : 45^\circ$ mixing)

for short times ($t \ll \tau$): $P_0(t) = \delta m^2 t^2$

for long times ($t \gg \tau$): $P_0(t) = \frac{1}{2}$

In medium ($\omega_B = \frac{1}{2} |\mu B| \gg \delta m$): $P_B(t) = \frac{\delta m^2}{\delta m^2 + \omega_B^2} \sin^2 \left(\sqrt{\delta m^2 + \omega_B^2} t \right),$

($\tau_B = \omega_B^{-1} \ll \tau, \sin^2 2\theta_B = \delta m^2 / \omega_B^2 \ll 1$)

for short times ($t \ll \tau_B$): $P_B(t) = \delta m^2 t^2$

for long times ($t \gg \tau_B$): $P_B(t) = \frac{1}{2} \frac{\delta m^2}{\omega_B^2}$

$$\Delta_B = P_0 - P_B > 0$$

Magnetic field **suppresses** oscillation. The experiments with the reactor neutrons in free flight as well in the UCN traps could observe the difference in the neutron lose rates for the magnetic field **on** and **off**

for more detailed discussion, see Pokotilovsky '06

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Neutron - Mirror neutron mixing in astrophysics

• *primordial baryon asymmetry can be generated via $\Delta B = 1$ processes like $udd \rightarrow u'd'd'$. The same (and possibly somewhat larger) baryon asymmetry would be generated in the Mirror sector, which could naturally explain the origin of the baryonic and dark matter balance in the Universe: $\Omega_D \sim \Omega_B$.*

N.B. *This mechanism does not require that $n - n'$ oscillation time should be necessarily small, within the present experimental reach. However, it requires that $\Delta B = 2$ processes like $udd \rightarrow \bar{u}\bar{d}\bar{d}$ should be also active though could be much slower. Hence, should the $n - n'$ oscillation be detected at the level $\tau_{nn'} < 10^4$ s, (i.e. $\mathcal{M}_{nn'} \sim 10$ TeV) it would give a strong argument that $n - \bar{n}$ oscillation should also exist at the experimentally accessible level, with the relevant cutoff scale $\mathcal{M}_{n\bar{n}} \sim 100$ TeV and thus $\tau_{n\bar{n}} \sim 10^9$ s.*

• *If $\tau_{nn'} < 10^3$ s, $n - n'$ oscillation provides an elegant mechanism for the transport of the ultra high energy cosmic rays at the large cosmological distances without suffering significant energy depression, and could be of interest in the search of the UHECR above the GZK cutoff and their correlation with the far distant astrophysical objects (BL Lacs, GRB's etc.)*

Z.B. & Bento '05

• *Fast $n - n'$ oscillation could have interesting implications also for the neutrons from the solar flares*

Mohapatra, Nasri, Nussinov '05

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Experimental limits & and future search

- ILL experiment for $n - \tilde{n}$ oscillation search in flight: $t \simeq 0.1$ s, $B < 10^{-4}$ G
 - no \tilde{n} event found, $\tau_{n\tilde{n}} > 10^8$ s (or > 3 yr) Baldo Ceolin et al. '94
 - as for $n - n'$: about 5% neutron deficit was observed, so taking

$$P_{nn'}(t) \simeq (t/\tau)^2 < 10^{-2}, \quad \tau_{nn'} > 1 \text{ s} \rightarrow \delta m < 10^{-15} \text{ eV}$$
- $n - n'$ – anomalous UCN losses, $\eta < 2 \cdot 10^{-6} \rightarrow \delta m < 3 \cdot 10^{-15} \text{ eV}$
- Nuclear Stability gives no limit for $\tau_{nn'}$ Z.B. & Bento '05

Recent Experimental search:

- $\tau > 2.7$ s Munich, Schmidt et al, Feb. 2007 (unpubl.)
- $\tau > 103$ s ILL Grenoble, Ban et al. May 2007, axXiv:0705.2336 [nucl-ex]
- $\tau > 414$ s ILL Grenoble, Serebrov et al. June 2007, axXiv:0706.3600 [nucl-ex]

Future experiments can reach sensitivity $\tau \sim 10^4$ s (DUSEL ??)

$n - n'$ oscillations can have very different experimental implications if n and n' states *are not exactly degenerate* at $B=0$. E.g. gravity is not *quite universal* between O and M matters, or there exist non-universal 5th forces of *non-gravitational* origin, or the *mirror magnetic field* is non-zero. *Opposite effect* is possible: magnetic field could *enhance the oscillation* instead of suppressing it.

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

$n - n'$ oscillation in mirror magnetic field ($B' \neq 0$)

Z. Berezhiani, arXiv: 0804.2088 [hep-ph]

Hamiltonian is 4×4 matrix describing oscillations and precessions

$$H_I = \begin{pmatrix} \mu(\vec{B} \cdot \vec{\sigma}) & \delta m \\ \delta m & \mu(\vec{B}' \cdot \vec{\sigma}) \end{pmatrix} = \begin{pmatrix} 2\vec{\omega} \cdot \vec{\sigma} & \delta m \\ \delta m & 2\vec{\rho} \cdot \vec{\sigma} \end{pmatrix}.$$

when oscillations can be averaged,

$$B = 0 : \quad P_0 = \frac{1}{2} \sin^2 2\theta_0 = \frac{\delta m^2}{2\rho^2} = 2 \left(\frac{\delta m}{\mu B'} \right)^2$$

$$B \neq 0 : \quad P(\vec{B}) = \frac{1}{2} \sin^2 2\theta = \frac{\delta m^2 (\vec{\rho} + \vec{\omega})^2}{2(\vec{\rho}^2 - \vec{\omega}^2)^2} = \frac{1 + \eta^2 + 2\eta \cos \beta}{(1 - \eta^2)^2} P_0$$

$\eta = B/B'$, β angle between \vec{B} and \vec{B}'


$$P_B = \frac{P(\vec{B}) + P(-\vec{B})}{2} = \frac{1 + \eta^2}{(1 - \eta^2)^2} P_0 \quad \longrightarrow \quad \Delta_B = P_B - P_0 = \frac{\eta^2(3 - \eta^2)}{(1 - \eta^2)^2} P_0$$

Δ_B is **positive** ($P_B > P_0$) if $\eta < \sqrt{3}$, i.e. $B' > 0.6B$

$$D_B(\beta) = \frac{P(\vec{B}) - P(-\vec{B})}{2} = \frac{2\eta}{(1 - \eta^2)^2} P_0 \cos \beta$$

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- **Oscillation**
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Comparison of observables



- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

$n - n'$ oscillation in mirror magnetic field

Experimental data from

G. Ban et al, PRL 99, 161603 (2007)

t_s [s]	50 (a)	50 (b)	100 (a)	175 (a)
$N_{B\uparrow}(t_*)$	44197 ± 53	44443 ± 53	28671 ± 30	17047 ± 31
$N_{B\downarrow}(t_*)$	44128 ± 53	44316 ± 46	28596 ± 30	16974 ± 31
$A_B(t_*) \times 10^3$	0.78 ± 0.85	1.43 ± 0.79	1.31 ± 0.74	2.15 ± 1.28
$N_0(t_*)$	44317 ± 40	44363 ± 53	28635 ± 21	17015 ± 22
$E_B(t_*) \times 10^3$	3.50 ± 1.24	-0.37 ± 1.43	0.05 ± 1.04	0.27 ± 1.83
κ_B	4.48 ± 5.12	-0.26 ± 1.00	0.04 ± 0.80	0.12 ± 0.85

Table 1: The UCN counts measured in configurations B_\uparrow, B_\downarrow ($B = 0.06$ G) and B_0 for different storage times t_s . Effective time $t_{\text{eff}} = t_s + (23 \pm 3)$ s.

$$D_B = (6.2 \pm 2.0) \times 10^{-7} \quad (\chi^2/d.o.f. = 0.52/3)$$

$$\Delta_B = (2.9 \pm 2.9) \times 10^{-7} \quad (\chi^2/d.o.f. = 6.9/3) \quad \text{Positive ?}$$

New Data (preliminary results) indicate 4.3 σ effect for D_B

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Spin dependent fifth forces

light pseudoscalar ϕ coupled with the nucleons of both sectors:

$$ig_p \phi (\bar{N} \gamma^5 N - \bar{N}' \gamma^5 N') = g_p \frac{\partial_\mu \phi}{2m} \cdot (\bar{N} \gamma^\mu \gamma^5 N - \bar{N}' \gamma^\mu \gamma^5 N')$$

Inhomogeneity of $\phi \rightarrow$ *spin-dependent forces* $\frac{\nabla \phi}{2m} \cdot (\bar{N} \Sigma N - \bar{N}' \Sigma N')$

If ϕ has also scalar couplings $g_s \phi (\bar{N} N + \bar{N}' N')$ Moody & Wilczek '84

then interaction potentials between two bodies are

(monopole)²: $V_{mm}(r) = -\frac{g_s^{(1)} g_s^{(2)}}{4\pi r} e^{-m_\phi r}$

monopole-dipole : $V_{md}(r) = \pm \frac{g_s^{(1)} g_p^{(2)} (\boldsymbol{\sigma} \cdot \boldsymbol{n})}{8\pi m_2} \left[\frac{m_\phi}{r} + \frac{1}{r^2} \right] e^{-m_\phi r}$

Therefore **the Earth** could be the source of $\nabla \phi$:
spin-dependent potential 13 orders of magnitude weaker than gravity would suffice

... but the source of $\nabla \phi$ could be some unknown matter with cosmological distribution

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- **Fifth Forces**
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Summary

Neutron - mirror neutron oscillation – violating **Baryon number** – can be rather fast, faster than neutron decay itself ... and it can have several intriguing implications for particle phenomenology (**underlying TeV physics at LHC**), astrophysics (**cosmic rays, instability of neutron stars, etc.**) and cosmology (**primordial baryogenesis: $\Omega'_B \geq \Omega_B$, BBN, etc.**).

May influence neutron lifetime measurements

$$\frac{N_{\text{fin}}}{N_{\text{in}}} = \exp[-\Gamma t + R\nu t + P_{nn'}\nu t] \quad P_{nn'}(\vec{B})$$

Easy to detect – comparing counts for different \vec{B} , regular loses **cancell out** . Crucial test: the neutron regeneration $n \rightarrow n' \rightarrow n$

Indications for “up-down” anomalies in the UCN counts with respect to \vec{B} direction – just statistical fluctuations ?

if not, can be explained by the mirror magnetic field $B = 0.03 - 3 \text{ G}$: on the Earth, in Solar System or in Galaxy

the latter \longrightarrow Day-Night (or siderial time) variations in comparing the UCN counts with opposite directions of the magnetic field.

would be good to check experimentally !

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- **Summary**
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Appendix 1: SUSY, GUT and O & M interactions

- Higgs-Higgs' quartic: $\lambda(\phi^\dagger\phi)(\phi'^\dagger\phi')$

could be interesting for the Higgs physics at the LHC ... but **BBN:** $\lambda < 10^{-8}$!

reaction $\phi\tilde{\phi} \rightarrow \phi'\tilde{\phi}'$ should not bring mirror sector in equilibrium with our particles

... natural in SUSY : lowest order mixed Higgs term is **D=5** operator

$$W = \frac{1}{M_{Pl}} (\phi_u\phi_d)(\phi'_u\phi'_d) \quad - \quad \text{in superpotential}$$

- Photon-photon' kinetic mixing: $\varepsilon F^{\mu\nu} F'_{\mu\nu}$

mirror particles become "millicharged" $Q' \sim \varepsilon Q$ relative to our photon Holdom '86

process $e^+e^- \rightarrow e'^+e'^-$ testable in positronium physics down to $\varepsilon \sim 10^{-7}$

Glashow '86, Gninenko '94

$N'N$ nuclear scattering testable in DM detectors down to $\varepsilon \sim 5 \cdot 10^{-9}$ Foot '03

but **BBN:** ($e^+e^- \rightarrow e'^+e'^-$ reaction) $\varepsilon < 3 \cdot 10^{-8}$ Carlson & Glashow '87

and **CMB+LSS:** $\varepsilon < 10^{-9}$ if mirror baryons constitute DM Z.B. & Lepidi '07

... natural in GUT : lowest order mixed gauge term is **D=6** operator

$$\mathcal{L} \sim \frac{\Sigma\Sigma'}{M_{Pl}^2} G^{\mu\nu} G'_{\mu\nu} \quad - \quad \text{e.g. } \Sigma^{(I)}, G_{\mu\nu}^{(I)} \sim 24\text{-plets in } SU(5) \times SU(5)'$$

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- **Appendix**
- Experiment
- Experiment
- Experiment
- Experiment

Search for $n - n'$ with the UCN storage

Counting of the UCN (ultra cold neutrons, velocities $v < 4 - 5$ m/s,) after a storage time t_s in a neutron trap, comparing the results for $B = 0$ and $B \neq 0$.

$$n(t_s) = n(t=0) \times \exp[-(\Gamma + \eta\nu + P_{nn'}(t_f)\nu) t_s]$$

t_f is a mean flight time between collisions ($\sim 0.05 - 0.1$ s),
 $\nu = 1/t_f$ is a collision frequency, and η is anomalous loss per collision by "natural" reasons (i.e. independent of magnetic field).

- For $B \neq 0$: $P_{nn'}(t_f) = \bar{P}_B \approx \frac{1}{2} \frac{\delta m^2}{\omega_B^2} = \frac{1}{2} \left(\frac{\tau_B}{\tau_0} \right)^2$

(Magnetic field is taken enough large to satisfy $t_f \gg \tau_B \approx \omega_B^{-1}$)

- For $B = 0$: $P_{nn'}(t_f) = \left(\frac{t_f}{\tau_0} \right)^2 \gg \bar{P}_B$

So signal is: $\frac{n(B=0, t_s)}{n(B, t_s)} = \exp[-a t_s] < 1$, i.e. a *should be positive*

Fitting $a = \frac{1}{t_f} \left(\frac{t_f}{\tau_0} \right)^2$ from the measurements, one finds $\tau_0 = \sqrt{t_f/a}$.

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D - Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Experiment Ban et al. May 2007

The UCN storage trap with volume 21 l, neutron velocities $v < 4$ m/s, wall collision frequency $\nu = 20 \text{ s}^{-1}$, average flight time $t_f = 0.05 \text{ s}$

"zero" magnetic field – $B_0 = (2 - 3) \cdot 10^{-5} \text{ G}$,

changing magnetic fields "up" and "down", $B_{\uparrow} = B_{\downarrow} = 0.06 \text{ G}$

storage times $t_s = 50, 100, 175 \text{ s}$, effective times $t^* = t_s + 23 \text{ s}$

● Expectation: $\frac{n(B=0, t^*)}{n(B_{\uparrow\downarrow}, t^*)} = \exp[-a t^*] < 1$, i.e. $a > 0$

● Fit of measurements: $a = -(5.4 \pm 5.8) \cdot 10^{-6} \text{ s}^{-1} \rightarrow \tau_0 > 103 \text{ s}$

$t^* \text{ [s]}$	73 (a)	73 (b)	123	198
$n(B_{\uparrow})$	44197 ± 53	44443 ± 53	28671 ± 30	17047 ± 31
$n(B=0)$	44317 ± 40	44363 ± 53	28635 ± 21	17015 ± 22
$n(B_{\downarrow})$	44128 ± 53	44316 ± 46	28596 ± 30	16974 ± 31
$n(B_{\uparrow\downarrow})$	44163 ± 38	44371 ± 35	28633 ± 22	17011 ± 22

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Up-Down Asymmetry

t^* [s]	$n(B=0)/n(B_{\uparrow\downarrow})$	$n(B_{\uparrow})/n(B_{\downarrow})$
73 (a)	1.0035 ± 0.0013	1.0016 ± 0.0017
73 (b)	0.9998 ± 0.0015	1.0028 ± 0.0016
73 (a + b)	1.0019 ± 0.0010	1.0022 ± 0.0012
123	1.0001 ± 0.0011	1.0026 ± 0.0015
198	1.0002 ± 0.0018	1.0043 ± 0.0026

● *Fit of* $\frac{n(B=0, t^*)}{n(B_{\uparrow\downarrow}, t^*)} = e^{\beta(t^*/t_f)} \approx 1 + \beta \left(\frac{t^*}{t_f} \right)$

$$\beta = (2.92 \pm 2.90) \times 10^{-7} \quad (68.27 \% CL)$$

● *Fit of* $\frac{n(B_{\uparrow}, t^*)}{n(B_{\downarrow}, t^*)} = 1 + \gamma \left(\frac{t^*}{t_f} \right)$

$$\gamma = (1.22 \pm 0.40) \times 10^{-6} \quad (68.27 \% CL)$$

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

$n - n'$ in non-degenerate case

$$H = \begin{pmatrix} m - i\Gamma/2 + V + \mu(\boldsymbol{\sigma} \cdot \mathbf{B}) & \delta m \\ \delta m & m' - i\Gamma'/2 + V' \end{pmatrix}$$

Consider $2\Delta E = (m' - m) + (V' - V) \neq 0$ – but small ($\sim 10^{-12}$ eV)

$$H_+ = \begin{pmatrix} m + V - 2\omega_B & \delta m \\ \delta m & m + V + 2\Delta E \end{pmatrix} \quad \text{for } \psi_+, \psi'_+ \text{ states,}$$

$$H_- = \begin{pmatrix} m + V + 2\omega_B & \delta m \\ \delta m & m + V + 2\Delta E \end{pmatrix} \quad \text{for } \psi_-, \psi'_- \text{ states}$$

Now (+) and (–) polarization states oscillate at different rates in magnetic medium, A being neutron polarization asymmetry:

$$\bar{P}_\pm(t) = \frac{1}{2} \frac{\delta m^2}{\delta m^2 + (\Delta E \pm \omega_B)^2}, \quad \bar{P}_0 = \frac{1}{2} \frac{\delta m^2}{\delta m^2 + \Delta E^2}, \quad (\omega_B = \frac{1}{2} |\mu B|)$$

● $\frac{n(B=0)}{n(B_{\uparrow\downarrow})}$ asymmetry $\beta = \left(\frac{\delta m}{\Delta E}\right)^2 \frac{\omega_B^2 (3\Delta E^2 - \omega_B^2)}{2(\Delta E^2 - \omega_B^2)^2} > 0$ if $\Delta E > 0.6\omega_B$

● $\frac{n(B_{\uparrow})}{n(B_{\downarrow})}$ asymmetry $\gamma = A \times \left(\frac{\delta m}{\Delta E}\right)^2 \frac{2\omega_B \Delta E^3}{(\Delta E^2 - \omega_B^2)^2}$ requires, $A > 30\%$

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

$n - n'$ in non-degenerate case

Consider now the case *but small* ($\sim 10^{-12}$ eV)

$$H_+ = \begin{pmatrix} m + V - 2\omega_B & \delta m \\ \delta m & m + V + 2\Delta E \end{pmatrix} \quad \text{for } \psi_+, \psi'_+ \text{ states ,}$$

$$H_- = \begin{pmatrix} m + V + 2\omega_B & \delta m \\ \delta m & m + V - 2\Delta E \end{pmatrix} \quad \text{for } \psi_-, \psi'_- \text{ states}$$

Now (+) and (−) polarization states oscillate at the same rates in magnetic medium, results do not depend on neutron polarization:

$$\bar{P}_\pm(t) = \frac{1}{2} \frac{\delta m^2}{\delta m^2 + (\Delta E \pm \omega_B)^2}, \quad \bar{P}_0 = \frac{1}{2} \frac{\delta m^2}{\delta m^2 + \Delta E^2}, \quad (\omega_B = \frac{1}{2} |\mu B|)$$

$$\bullet \frac{n(B=0)}{n(B_{\uparrow\downarrow})} \text{ asymmetry } \beta = \left(\frac{\delta m}{\Delta E}\right)^2 \frac{\omega_B^2 (3\Delta E^2 - \omega_B^2)}{2(\Delta E^2 - \omega_B^2)^2} > 0 \text{ if } \Delta E > 0.6\omega_B$$

$$\bullet \frac{n(B_{\uparrow})}{n(B_{\downarrow})} \text{ asymmetry } \gamma = \left(\frac{\delta m}{\Delta E}\right)^2 \frac{2\omega_B \Delta E^3}{(\Delta E^2 - \omega_B^2)^2}$$

one can fit both data for $\Delta E \sim 10^{-12}$ eV

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Mirror Physics: Summary

- String Theory: parallel D-branes or brane-antibrane
- restoring Parity: $L \leftrightarrow R$ – can remain exact (models of exact mirror parity) or spontaneously broken Z.B., Dolgov & Mohapatra 96
- Common gauge forces between two sectors: e.g. $U(1)_{B-L}$, or (anomaly free) gauge flavour symmetry $SU(3)_H$ between fermion families: helps for SUSY flavour changing problem (D-terms) Z.B., 98
- Higgs sector stability: Higgs as pseudoGoldstone in SUSY – accidental global $U(4)$ symmetry Z.B., 05
Falkowski, Pokorski & Schmalz 06
- Photon-photon' kynetic mixing (invisible 0- P_s decay) Holdom, Glashow
neutrino-neutrino' (active - sterile) mixing Foot & Volkas; ZB & Mohapatra
neutron - neutron' mixing (hydrogen - hydrogen') mixing ZB & Bento
pion - pion' mixing (DAMA vs. CDMS) ZB, Panci & Rossi
Kaon - Kaon' mixing (new features in CP-violation ?) etc. etc.
- Strong CP-problem: new models for axion avoiding mass-coupling correlation $m_a \sim f_\pi m_{pi} / f_a$ Z.B., Gianfagna, Gianotti 2000

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment


Mirror Cosmology & Astrophysics: Summary

- Mirror sector should be cooler than ours: $T'/T < 0.5$ or so (BBN)
- Dark matter of the Universe:
self-interacting dissipative (for exact parity): requires $T'/T < 0.2$
or WDM, or almost CDM (for broken parity)
- Baryogenesis & dark matter genesis: Bento & Z.B., 2001
Understanding why $\Omega_D \sim \Omega_B$
- Microlensing (MACHOs) Z.B., Dolgov & Mohapatra 96, Blinikov 98
- Gamma- Ray Bursts and Supernove Blinnikov 98; Z.B. & Drago 99
- Super high energy neutrinos Berezinsky & Vilenkin, 2000
– Propagation of ultra High energy protons Z.B. & Bento 05
- Quasars & supermassive black holes ZB, Comelli & Villante 2000
- Possible dark matter detection (DAMA vs. others) Foot 2003
- Invisible planets and meteorits (Tunguska) Foot, Silagadze
Thermal imprints of mirror matter Foot & Mitra

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Mirror Matter Evolution epochs

Z.B., Comelli & Villante, '01



- Earlier decoupling of mirror photons ($x < 0.5$):



$$z'_{\text{dec}} \simeq x^{-1} z_{\text{dec}} \quad x_{\text{eq}} = 0.05(\Omega_M h^2)^{-1} \simeq 0.3$$

- Jeans scale is smaller : $\lambda_J \ll d_H$ before Matter=Radiation

- Silk scale is smaller : $\lambda'_S \sim 5x_{\text{eq}}^{5/4} (x/x_{\text{eq}})^{3/2} (\Omega_M h^2)^{-3/4}$ Mpc

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment


CMB & LSS power spectra



- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment
- Experiment

LSS power spectra

Z.B., Ciarcelluti, Comelli & Villante, '03



- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Cosmic Fine Tuning?

The early Universe:

- multi-stage: Inflation \rightarrow (re)heating \rightarrow Friedmann epoch ...
- Universe is flat and homogeneous ...
- Adiabatic perturbations with nearly flat spectrum ...

Today's Universe:

- multi-component: visible matter, dark matter, dark energy ...
- $\Omega_{\text{tot}} \approx 1$ Universe is flat: $\rho_{\text{tot}} = \rho_{\text{cr}}$...
- $\Omega_{\text{B}} \simeq 0.04$ visible (Baryon) matter is a small fraction ...
- $\Omega_{\text{D}} \simeq 0.20$ dark matter: **WIMPS? Axions?**
- $\Omega_{\Lambda} \simeq 0.75$ dark energy: **Λ -term? 5th-essence?**

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Cosmic Fine Tuning?

The early Universe:

- multi-stage: Inflation → (re)heating → Friedmann epoch ...
- Universe is flat and homogeneous ...
- Adiabatic perturbations with nearly flat spectrum ...

Today's Universe:

- multi-component: visible matter, dark matter, dark energy ...
- $\Omega_{\text{tot}} \approx 1$ Universe is flat: $\rho_{\text{tot}} = \rho_{\text{cr}}$...
- $\Omega_{\text{B}} \simeq 0.04$ visible (Baryon) matter is a small fraction ...
- $\Omega_{\text{D}} \simeq 0.20$ dark matter: **WIMPS? Axions?**
- $\Omega_{\Lambda} \simeq 0.75$ dark energy: **Λ -term? 5th-essence?**

Anthropic World? Origin and nature of DM & DE remain open!

Unified picture for BM and DM – Mirror Matter?

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Boltzmann Eqs.

Evolution for (B-L)' and (B-L) $T_R \ll M$

$$\frac{dn_{B-L}}{dt} + 3Hn_{B-L} + \Gamma n_{B-L} = \frac{3}{4} \Delta\sigma n_{\text{eq}}^2$$

$$\frac{dn'_{B-L}}{dt} + 3Hn'_{B-L} + \Gamma' n'_{B-L} = \frac{3}{4} \Delta\sigma' n_{\text{eq}}^2$$

$\Gamma \propto n'_{\text{eq}}/M^2$ is the effective reaction rate of $\Delta L' = 1$ and $\Delta L' = 2$ processes

$$\Gamma'/\Gamma \simeq n'_{\text{eq}}/n_{\text{eq}} \simeq x^3 ; \quad x = T'/T$$

$$\Delta\sigma' = -\Delta\sigma = \frac{3\varepsilon_{CP} S}{32\pi^2 M^4}$$

where $S \sim 16T^2$ is the c.m. energy square,


$$\varepsilon_{CP} = \text{Im Tr}[(y^\dagger y)^* g^{-1} (y'^\dagger y') g^{-2} (y^\dagger y) g^{-1}]$$

$$Y_{BL} = D(k) \cdot Y_{BL}^{(0)} ; \quad Y'_{BL} = D(kx^3) \cdot Y_{BL}^{(0)}$$

$$Y_{BL}^{(0)} \approx 2 \times 10^{-3} \frac{\varepsilon_{CP} M_{Pl} T_R^3}{g_*^{3/2} M^4} .$$

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Exact M-parity: $M'_N = M_N$



$$n_B/n'_B = D(k), \quad k = [\Gamma_{\text{eff}}/H]_{T=T_R} : \quad \Omega_B/\Omega'_B \simeq 0.15 - 1$$

Depletion factor $D(k) = \frac{3}{5} e^{-k} F(k) + \frac{2}{5} G(k)$; for $k \ll 1$, $D(k) = 1$

$$F(k) = \frac{1}{4k^4} [(2k-1)^3 + 6k - 5 + 6e^{-2k}] : \quad T > T_R,$$

$$G(k) = \frac{3}{k^3} [2 - (k^2 + 2k + 2)e^{-k}] : \quad T < T_R$$


$$\text{Heating: } \Delta N_\nu \simeq k/g_* \quad x = (k/6g_*)^{1/4} < 0.2 : \quad k \leq 2, \quad (\text{LSS})$$

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment

Broken M parity: $M'_W > M_W$?

Spont. broken M parity: $v' \gg v$

Z.B., Dolgov & Mohapatra '96



$$n'_B \simeq n_B \quad k < 1 \text{ (robust non-equilibrium)}$$

$M'_N/M_N \simeq (\Lambda'/\Lambda)$ changes slowly with M'_W

$m'_e/m_e \simeq M'_W/M_W$ changes fastly with M_W .

– Properties of MB's get closer to CDM : $M'_W \sim 10 \text{ TeV}$?

- Carrol's Alice...
- Mirror World
- Mirror Particles
- Interactions
- B & L violation
- BBN demands
- Present Cosmology
- Visible vs. Dark matter
- B vs. D – Fine Tuning demonstration
- Unification
- Neutrino Mixing
- See-Saw
- Leptogenesis: diagrams
- Leptogenesis: formulas
- Epochs
- Neutron mixing
- Neutron mixing
- Oscillation
- Probabilities
- Neutron mixing
- Experiment
- Oscillation
- Oscillation
- Oscillation
- Fifth Forces
- Summary
- Appendix
- Experiment
- Experiment
- Experiment
- Experiment