


WISPy Cold Dark Matter



J. Jaeckel^{**}

S. Abel[†], J. Berges^{**}, B. Doebrich^z, M. Goodsell^{xx},
H. Gies^o, F. Kahlhoefer^{*}, S. Knirck^{**},
V. Khoze[†], A. Lobanov^y, J. Redondo^x,
A. Ringwald^{*}, K. Schmidt-Hoberg^{*}
and The FUNK Collaboration

^{**}ITP Heidelberg, ^zCERN, [†]IPPP Durham, ^{*}DESY,
^yMPIfR Bonn, ^xU. Zaragoza, ^{xx}Paris LPTHE, ^oITP Jena




$$-\mathcal{L} \supset \frac{1}{2}m_h^2 h^2 + \sqrt{\frac{\eta}{2}}m_h h^3 + \frac{\eta}{4}h^4$$
$$-g_v m_V V^2 h + \frac{m_f \bar{f} f}{v}$$
$$-\frac{1}{4} \frac{\alpha_s}{12\pi} G_{\mu\nu}^a G^{a\mu\nu} \log(1 + \frac{h}{v})$$

+ nothing else


What we know...

Inventory of the Universe




Where does it hide?

CALVIN and HOBBS BY WATSON




Where does it hide?

CALVIN and HOBBS BY WATSON



Exploring is (at least) 2 dimensional




Can Dark Matter be WISPy?

(Weakly Interacting Sub-eV Particley)
Slim




Properties of Dark Matter

- Dark matter is dark, i.e.
it doesn't radiate!
(and also doesn't absorb)
- very, very weak interactions with light
and with ordinary matter
- Exactly the property of
WISPs



Exploring is (at least) 2 dimensional




A common prejudice

- Dark Matter has to be heavy: $m_{\text{DM}} \gtrsim \text{keV}$.
- Prejudice based on thermal production!
and/or fermionic DM!

Both assumptions give
minimal velocity

→ galaxy,
i.e. structure,
formation inhibited!



Weakly interacting sub-eV DM

- Has to be non-thermally (cold!!!) produced

→ See misalignment mechanism ✓

- Bosonic!

→ Axion(-like particles)
Hidden Photons ✓

Dark matter has to be heavy...

Dark matter has to be heavy $m_{\text{DM}} \gtrsim \text{keV}$?

Dark matter has to be heavy...



Example WISP:

Axions in a nutshell


A dirty little secret...

$$S = \int d^4x \left[-\frac{1}{4} G^{\mu\nu} G_{\mu\nu} - \frac{\theta}{4} G^{\mu\nu} \tilde{G}_{\mu\nu} + i\bar{\psi} D_\mu \gamma^\mu \psi + \bar{\psi} M \psi \right]$$

" $\sim \theta \vec{E} \cdot \vec{B}$ "

- The θ -term violates time reversal (T=CP)!
- Connected to strong interactions!


 Electric dipole moment
of the neutron!



Not found
 $\Rightarrow \theta \sim 0!!!$


The axion solution to the strong CP problem

- Make θ dynamical \rightarrow it can change its value




The axion solution to the strong CP problem

- Make θ dynamical \rightarrow it can change its value



The axion solution to the strong CP problem


- Make θ dynamical \rightarrow it can change its value



\rightarrow QCD likes to be CP conserving (if we allow it)

The axion solution to the strong CP problem

- Make θ dynamical \rightarrow it can change its value



\rightarrow Can still move

\rightarrow new particle = axion


- Classical flatness from symmetry
- Quantum corrections are small
- New **light** particle: **The Axion**
(it's a **Weakly Interacting Sub-eV Particle**)

Dark matter candidate

**Good motivation
for axion/WISP experiments**

The diagram consists of three yellow arrows. One arrow points from the word 'light' in the third bullet point to the text 'Dark matter candidate'. A second arrow points from 'Dark matter candidate' to the text 'Good motivation for axion/WISP experiments'. A third arrow points from the word 'light' in the third bullet point directly to 'Good motivation for axion/WISP experiments'.

Axion also couples to two photons




$$\mathcal{L} = -\frac{1}{4}F^{\mu\nu}F_{\mu\nu} + \frac{1}{2}\partial_\mu a\partial^\mu a - m^2 a^2 - \frac{1}{4}g_{a\gamma\gamma}aF^{\mu\nu}\tilde{F}_{\mu\nu} + \dots$$

Coupling to two photons


Very very weak $g_{a\gamma\gamma} \sim \frac{\alpha}{2\pi f_a}$

Because: Very large

Axion-like Particles




Axion-like Particles



Looking for Axions/ALPs


Light shining through walls

“Light shining through a wall”



Light shining through walls

“Light shining through a wall”




- **Test** $P_{\gamma \rightarrow X \rightarrow \gamma} \lesssim 10^{-20}$
- **Enormous precision!**
- **Study extremely weak couplings!**

Photons coming through the wall!

- It could be Axion(-like particle)s!

- Coupling to two photons: $\frac{1}{M} a \tilde{F} F \sim \frac{1}{M} a \vec{E} \cdot \vec{B}$




$$P_{\gamma \rightarrow a \rightarrow \gamma} \sim N_{\text{pass}} \left(\frac{BL}{M} \right)^4$$

ALPS @ Hamburg

$\sim 10^4$ GeV

$\sim 10^8$ GeV




Going deeper

Helioscopes


CAST@CERN

SUMICO@Tokyo

SHIPS@Hamburg




“Light shining through a wall”




Going to the future: IAXO

The International Axion Observatory




CAST + IAXO

$\sim 10^9$ GeV




Dark Matter(s)


The axion has no clue where to start



The axion has no clue where to start




The axion solution to the strong CP problem



- Oscillations contain energy
- behave like non-relativistic particles ($T=0$)

Why Cold? Inflation!

Field
value




space

$$velocity \sim \frac{p}{m} \sim \frac{\hbar}{m} \frac{d}{dx} \rightarrow 0$$

Axion(-like particle) Dark Matter

$\sim 10^7 \text{ GeV}$


$\sim 10^{12} \text{ GeV}$




Detecting WISPy DM

Use a plentiful source of axions


- Photon Regeneration



Signal: Total energy of axion




$$h\nu = m_a c^2 [1 + \mathcal{O}(\beta^2 \sim 10^{-6})]$$



Virial velocity
in galaxy halo!


An extremely sensitive probe!!!



Electricity from Dark Matter ;-).


- Photon Regeneration

Photon
(amplified in resonator)



axion

(dark matter)



10^{-23}W!



Really sustainable Energy

- Galaxy contains $(6-30)\times 10^{11}$ solar masses of DM

→ $(3-15)\times 10^{43}$ TWh

@100000 TWh per year (total world today)


→ 10^{38} years ☺

DM power


$$\rho * v \sim 300 \text{ MeV/cm}^3 * 300 \text{ km/s} \sim 10 \text{ W/m}^2$$

compared to 2 W/m^2 for wind

A discovery possible any minute!




Encircling the axion...






Beyond ALPs

Also for hidden photons!!!

- There are other very light DM candidates
- E.g
extra (hidden) U(1) bosons=hidden photons!!!



@ DESY + Bonn: WISPDMMX



New Results!

1. WISPDMMX: A haloscope for WISP Dark Matter between 0.8-2 μeV


Le Hoang Nguyen, Dieter Horns, Andrei Lobanov, Andreas Ringwald. Nov 10, 2015.

DESY-15-185


e-Print: [arXiv:1511.03161](https://arxiv.org/abs/1511.03161) [physics.ins-det] | [PDF](#)

Broadband Search Strategy

Dark Matter Antenna



Probes here;
very sensitive!!




The FUNK Experiment

Recycle Auger mirror


Detector



First Results





Upgrade: The PMT 9000(+107)





Discovery Potential 😊!!!

A Dream for Astrology ehm Astronomy

- Emission from moving dark matter



- A picture of the DM-velocity distribution




Going
Monodromic

Axion Monodromy

- Allows for extended field range

$$V(\phi) = \frac{1}{2}m^2\phi^2 + \Lambda^4 \left(1 - \cos \left(\frac{\phi}{2\pi f} \right) \right)$$




Advantages



- Allows to start with higher energy density
→ More DM




VS




Models
in this region!



Interesting Phenomena??




Could get stuck here




Oscillations like DM!

Interesting Phenomena??



Interesting Phenomena??




Regions with "negative mass"


Instability \rightarrow Particle Production with $p \neq 0$?!?

Running through walls


A WISPy Domain Wall




Domain wall
from side 1
To side 2



A WISPy Domain Wall




Domain wall
from side 1
To side 2



aLIGO

- Has detected gravitational waves!!
- Is an Interferometer



Interference pattern


Causing a phase shift

- Interaction inside wall creates photon mass

$$\mathcal{L}_A = -\frac{1}{4}F^{\mu\nu}F_{\mu\nu} - \frac{1}{2}m_{0,\gamma}^2 \sin^2 \left(\frac{N_A \phi}{f} \right) A^\mu A_\mu$$

aLIGO

- Has detected gravitational waves!!
- Is an Interferometer



— Interference pattern **changed** —

Signal shapes




FIG. 6: $L = 4000$ m, $\omega \approx 1$ eV, $m = 10$ neV, $m_{\gamma,0} = 1$ neV, $N_A/N_\phi = 1$, $\alpha = \pi/2.2, \pi/2.5, \pi/3$ (black, blue, red), v chosen such that signal has roughly a length of $0.02s \sim 1/(50$ Hz) this corresponds to $v = 1 \times 10^{-3}$.

Signal shapes




FIG. 8: As in Fig. ?? but $m_{\gamma,0} = 0.1 \text{ neV}$, $N_A/N_\phi = 5$, $m = 0.5 \text{ neV}$, $\alpha = \pi/2$ and $v = 1 \times 10^{-3}$.

How to distinguish from grav waves?

- $velocity \ll c$
 - $v \sim 10^{-3}$
-
- Time difference between two sites
 ~few seconds
 - Need careful analysis strategies
-


Conclusions

Conclusions


- Good Physics Case for Axions and WISPs

➔ explore 'The Low Energy Frontier'

- Low energy experiments complementary to accelerators!



- Dark Matter may be WISPy 😊
 - ➔ New Search opportunities!
 - ➔ Searches ongoing!
 - ➔ Crazy things to explore!



Hidden sector

